

Chapter 147

Non-Inferiority Tests for the Difference of Two Within-Subject CV's in a Parallel Design

Introduction

This procedure calculates power and sample size of non-inferiority tests for the difference of within-subject coefficients of variation (CV) from a parallel design with replicates (repeated measurements) of a particular treatment. This routine deals with the case in which the statistical hypotheses are expressed in terms of the difference of the within-subject CVs, which is the standard deviation divided by the mean.

Technical Details

This procedure uses the formulation first given by Quan and Shih (1996). The sample size formulas are given in Chow, Shao, Wang, and Lokhnygina (2018).

Suppose x_{ijk} is the response in the i th group or treatment ($i = 1, 2$), j th subject ($j = 1, \dots, N_i$), and k th measurement ($k = 1, \dots, M$). The simple one-way random mixed effects model leads to the following estimates of CV1 and CV2

$$\widehat{CV}_i = \frac{\hat{\sigma}_i}{\hat{\mu}_i}$$

$$\hat{\mu}_i = \frac{1}{N_i M} \sum_{j=1}^{N_i} \sum_{k=1}^M x_{ijk}$$

$$\hat{\sigma}_i^2 = \frac{1}{N_i(M-1)} \sum_{j=1}^{N_i} \sum_{k=1}^M (x_{ijk} - \bar{x}_{ij\cdot})^2$$

where

$$\bar{x}_{ij\cdot} = \frac{1}{M} \sum_{k=1}^M x_{ijk}$$

Testing Non-Inferiority

The following hypotheses are usually used to test for the non-inferiority of CV

$$H_0: CV_1 - CV_2 \geq D0 \quad \text{versus} \quad H_1: CV_1 - CV_2 < D0.$$

The one-sided test statistic used to test this hypothesis is

$$T = \frac{(\widehat{CV}_1 - \widehat{CV}_2) - D0}{\sqrt{\frac{\hat{\sigma}_1^{*2}}{N_1} + \frac{\hat{\sigma}_2^{*2}}{N_2}}}$$

where $D0$ is the hypothesized CV difference under the null hypothesis and

$$\hat{\sigma}_i^{*2} = \frac{1}{2M} \widehat{CV}_i^2 + \widehat{CV}_i^4$$

T is asymptotically distributed as a standard normal random variable.

Hence the null hypothesis is rejected if $T < z_\alpha$.

Power

The power of this combination of tests is given by

$$\text{Power} = \Phi(z_\alpha - \mu_z)$$

where

$$\sigma_i^{*2} = \frac{1}{2M} CV_i^2 + CV_i^4$$

$$\mu_z = \frac{(CV_1 - CV_2) - D0}{\sqrt{\frac{\sigma_1^{*2}}{N_1} + \frac{\sigma_2^{*2}}{N_2}}}$$

and $\Phi(x)$ is the standard normal CDF.

A simple binary search algorithm can be applied to this power function to obtain an estimate of the necessary sample size.

Example 1 – Finding Sample Size

A company has developed a generic drug for treating rheumatism and wants to show that its within-subject CV is non-inferior to a reference drug. A parallel design with 2 repeated measurements per subject will be used.

Company researchers set the significance level to 0.05, the power to 0.90, CV2 to 0.4, CV1.0 to 0.5, and CV1.1 to 0.2 0.3 0.4 0.45. They want to investigate the range of required sample size values assuming that the two group sample sizes are equal.

Setup

If the procedure window is not already open, use the PASS Home window to open it. The parameters for this example are listed below and are stored in the **Example 1** settings file. To load these settings to the procedure window, click **Open Example Settings File** in the Help Center or File menu.

Design Tab	
Solve For	Sample Size
Power.....	0.90
Alpha.....	0.05
Group Allocation	Equal (N1 = N2)
M (Measurements Per Subject)	2
Input Type.....	Differences
D0 (Non-Inferiority Difference)	0.1
D1 (Actual Difference).....	0.05 0 -0.05 -0.1
CV2 (Group 2 Coef of Variation).....	0.4

Non-Inferiority Tests for the Difference of Two Within-Subject CV's in a Parallel Design

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Reports

Numeric Results

Solve For: [Sample Size](#)

Hypotheses: $H_0: CV1 - CV2 \geq D_0$ vs. $H_1: CV1 - CV2 < D_0$ (One-Sided)

Power		Sample Size				Coefficient of Variation					
						Measurements per Subject M	Non- Inferiority CV1.0	Actual CV1.1	Reference CV2	Difference	
										Non- Inferiority D0	Actual D1
Target	Actual	N1	N2	N							Alpha
0.9	0.9049	21	21	42	2	0.5	0.30	0.4	0.1	-0.10	0.05
0.9	0.9040	43	43	86	2	0.5	0.35	0.4	0.1	-0.05	0.05
0.9	0.9015	113	113	226	2	0.5	0.40	0.4	0.1	0.00	0.05
0.9	0.9002	539	539	1078	2	0.5	0.45	0.4	0.1	0.05	0.05

Target Power	The desired power value entered in the procedure. Power is the probability of rejecting a false null hypothesis.
Actual Power	The actual power achieved. Because N1 and N2 are discrete, this value is usually slightly larger than the target power.
N1	The number of subjects from group 1. Each subject is measured M times.
N2	The number of subjects from group 2. Each subject is measured M times.
N	The total number of subjects. $N = N1 + N2$.
M	The number of measurements per subject.
CV1.0	The non-inferiority boundary. CVs below this value are concluded as non-inferior.
CV1.1	The actual CV of group 1 at which the power is calculated (the value of CV1 assumed by H1).
CV2	The within-subject coefficient of variation in group 2 assumed by both H0 and H1.
D0	The non-inferiority difference (CV1.0 - CV2). $D0 = CV1.0 - CV2$.
D1	The actual difference (CV1.1 - CV2) at which the power is calculated (assumed by H1). $D1 = CV1.1 - CV2$.
Alpha	The probability of rejecting a true null hypothesis.

Summary Statements

A parallel two-group design with replicates will be used to test whether the Group 1 coefficient of variation (σ_1 / μ_1) is non-inferior to the Group 2 coefficient of variation (σ_2 / μ_2), by testing whether the difference in within-subject coefficients of variation is less than 0.1 ($H_0: CV1 - CV2 \geq 0.1$ versus $H_1: CV1 - CV2 < 0.1$). Each subject will be measured 2 times. The comparison will be made using a one-sided, two-sample Z-test with a Type I error rate (α) of 0.05. To detect a within-subject coefficient of variation difference of -0.1 (CV1 = 0.3, CV2 = 0.4) with 90% power, the number of subjects needed will be 21 in Group 1, and 21 in Group 2.

Non-Inferiority Tests for the Difference of Two Within-Subject CV's in a Parallel Design

Dropout-Inflated Sample Size

Dropout Rate	Sample Size			Dropout-Inflated Enrollment Sample Size			Expected Number of Dropouts		
	N1	N2	N	N1'	N2'	N'	D1	D2	D
20%	21	21	42	27	27	54	6	6	12
20%	43	43	86	54	54	108	11	11	22
20%	113	113	226	142	142	284	29	29	58
20%	539	539	1078	674	674	1348	135	135	270

Dropout Rate	The percentage of subjects (or items) that are expected to be lost at random during the course of the study and for whom no response data will be collected (i.e., will be treated as "missing"). Abbreviated as DR.
N1, N2, and N	The evaluable sample sizes at which power is computed. If N1 and N2 subjects are evaluated out of the N1' and N2' subjects that are enrolled in the study, the design will achieve the stated power.
N1', N2', and N'	The number of subjects that should be enrolled in the study in order to obtain N1, N2, and N evaluable subjects, based on the assumed dropout rate. After solving for N1 and N2, N1' and N2' are calculated by inflating N1 and N2 using the formulas $N1' = N1 / (1 - DR)$ and $N2' = N2 / (1 - DR)$, with N1' and N2' always rounded up. (See Julious, S.A. (2010) pages 52-53, or Chow, S.C., Shao, J., Wang, H., and Lohknygina, Y. (2018) pages 32-33.)
D1, D2, and D	The expected number of dropouts. $D1 = N1' - N1$, $D2 = N2' - N2$, and $D = D1 + D2$.

Dropout Summary Statements

Anticipating a 20% dropout rate, 27 subjects should be enrolled in Group 1, and 27 in Group 2, to obtain final group sample sizes of 21 and 21, respectively.

References


- Quan, H. and Shih, W.J. 1996. 'Assessing reproducibility by the within-subject coefficient of variation with random effects models'. Biometrics, 52, pages 1195-1203.
- Chow, S.C., Shao, J., Wang, H., and Lohknygina, Y. 2018. Sample Size Calculations in Clinical Research, Third Edition. Taylor & Francis/CRC. Boca Raton, Florida.

This report gives the sample sizes for the indicated scenarios.

Non-Inferiority Tests for the Difference of Two Within-Subject CV's in a Parallel Design

Plots Section

Plots



This plot shows the relationship between sample size and D1.

Example 2 – Validation using Chow et al. (2018)

Chow *et al.* (2018) pages 203-204 presents an example of a one-sided, lower-tail test in which $CV1.1 = 0.5$, $CV1.0 = 0.8$, $CV2 = 0.7$, $M = 2$, $\alpha = 0.05$, and power = 0.8. The sample size is found to be 34 per group.

Setup

If the procedure window is not already open, use the PASS Home window to open it. The parameters for this example are listed below and are stored in the **Example 2** settings file. To load these settings to the procedure window, click **Open Example Settings File** in the Help Center or File menu.

Design Tab

Solve For **Sample Size**
 Power..... **0.80**
 Alpha..... **0.05**
 Group Allocation **Equal (N1 = N2)**
 M (Measurements Per Subject) **2**
 Input Type..... **Coefficients of Variation**
 CV1.0 (Non-Inferiority Coef of Variation) **0.8**
 CV1.1 (Actual Coef of Variation)..... **0.5**
 CV2 (Group 2 Coef of Variation)..... **0.7**

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results

Solve For: [Sample Size](#)

Hypotheses: $H_0: CV1 - CV2 \geq D_0$ vs. $H_1: CV1 - CV2 < D_0$ (One-Sided)

					Coefficient of Variation						
					Measurements per Subject M	Difference					Alpha
Power		Sample Size				Non- Inferiority CV1.0	Actual CV1.1	Reference CV2	Non- Inferiority D0	Actual D1	
Target	Actual	N1	N2	N							
0.8	0.8052	34	34	68	2	0.8	0.5	0.7	0.1	-0.2	

The sample sizes match Chow et al. (2018).