Bridging Study Test of Sensitivity using a

Two-Group T-Test (Continuous Outcome)

Introduction

This procedure calculates the power and sample size required for bridging studies that two-sample, equalvariance t-test to estimate the sensitivity of the anticipated bridging study. The response data is continuous (assumed normal). Only a brief introduction to the subject will be given here. For a comprehensive discussion, refer to Chow, Shao, and Hu (2002).

Bridging Studies

Once a pharmaceutical product has been approved for use in one or more regions (countries) through a set of clinical trials, it is often desirable to register the product in a new region that was not included in the original study. When the cost and time needed to complete an additional set of clinical trials in the new region is prohibitive, a bridging methodology may be used to obtain the approval. After a value of the sensitivity index is obtained, an assessment must be made to determine whether the bridging study is necessary. If the bridging study is required, further analysis must be made to determine how large of a sample size is needed. This procedure helps with both of these questions.

Sensitivity Index

Suppose a randomized, parallel-group, placebo-controlled study has been completed and used to evaluate a test compound for use in treating a certain disease in a particular population. Now suppose that the sponsors of the test compound want to obtain approval for use in another population. This new population may be a demographic (e.g., elderly) that was not represented in the original study, or it may be an additional world region (e.g., Japan). It is assumed that the response to the test compound will be similar but slightly different in the new region. The similarities between the response distributions in the two regions can be summarized by their means and variances.

As is described below, Chow, Shao, and Hu (2002) proposed that these population differences could be summarized as a single number called the *sensitivity index*. They proposed that this index be used when there are well known ethnic differences between the two populations. Rather than have two parameters (means and standard deviations) to use for evaluating the population differences, they proposed that the ratio of each region's signal-to-noise ratios could be used.

Test Procedure for Assessing Sensitivity

Suppose a clinical trial using a typical parallel-group design has been completed and the treatment has been found to be beneficial in treating a certain disease. The analysis of this study, which is called the *original study*, is summarized as follows.

Let X_{ij} be the response of subject *j* on receiving treatment *i* in original study where *i* = T (treatment), C (control) and $j = 1, ..., N_i$. Let $N = N_T + N_C$. Assume that the X_{ij} are independently normally distributed with means μ_i and common variance σ^2 . Suppose the statistical hypotheses to be tested are

$$H_0: \mu_T - \mu_C = 0$$
 vs. $H_1: \mu_T - \mu_C \neq 0$

Assuming that the standard two-sample t-test is used for the analysis. The test statistic

$$T = \frac{\bar{x}_T - \bar{x}_C}{\sqrt{\left(\frac{(N_T - 1)S_T^2 + (N_C - 1)S_C^2}{N - 2}\right)\left(\frac{1}{N_T} + \frac{1}{N_C}\right)}}$$

is assumed to be distributed as a *Student's t* random variable with the usual degrees of freedom. The power of *T* is

$$p(\theta) = 1 - T_{N-2}'\left(t_{N-2,1-\frac{\alpha}{2}} \middle| \theta\right) + T_{N-2}'\left(-t_{N-2,1-\frac{\alpha}{2}} \middle| \theta\right)$$

where $t_{N-2,1-\frac{\alpha}{2}}$ is t-value from the central *t* distribution, T'_{N-2} is the distribution function of the noncentral *t* distribution, and θ is the noncentrality parameter calculated as

$$\theta = \frac{\mu_T - \mu_C}{\sigma_{\sqrt{\left(\frac{1}{N_T} + \frac{1}{N_C}\right)}}}$$

Suppose *x* represents the data from the original trial and T(x) is the value of *T* based on *x*. If θ is replaced by T(x) in the power function above, the *estimated power* $\hat{P}(T(x)) = p(T(x))$ is defined by Shao and Chow (2002) as the *reproducibility probability* for the bridging trial with the same subject population and with the same sample size.

Define T^* to be the value of T obtained when N_T and N_C of the original study is replaced by N_T^* and N_C^* which are the corresponding sample sizes of the bridging study. The estimated power of the new study is given by $\hat{P}(T^*(x))$.

Suppose the mean responses in the treatment and control groups of the original study are named μ_{OT} and μ_{OC} . Further suppose a single variance, σ_O^2 , can be used for both groups of the original study. Similar quantities in the new region are μ_{BT} , μ_{BC} , and σ_B^2 .

Let the relative difference between means in the two regions be defined as

$$R_{MD} = \frac{(\mu_{BT} - \mu_{BC}) - (\mu_{OT} - \mu_{OC})}{(\mu_{OT} - \mu_{OC})}$$

Similarly, let the ratio between standard deviations in the two regions be defined as

$$R_{SD} = \frac{\sigma_B}{\sigma_O}$$

The signal-to-noise ratio of the difference in the bridging study is $(\mu_{OT} - \mu_{OC})/\sigma_0$. Similarly, the signal-to-noise ratio of the difference in the bridging study is $(\mu_{BT} - \mu_{BC})/\sigma_B$.

The ratio of the two-study region signal-to-noise ratios of the differences between these parameters is

$$\Delta = \frac{(\mu_{BT} - \mu_{BC})/\sigma_B}{(\mu_{OT} - \mu_{OC})/\sigma_0}$$

which is called the sensitivity index. A little algebra will show that this can be rewritten as

$$\Delta = \frac{1 + R_{MD}}{R_{SD}}$$

Hence Δ is a measure of the change in signal-to-noise ratio from the original study to the bridging study.

Note that Δ may be computed for different values of R_{MD} and R_{SD} using the *Bridging Study Sensitivity Index* procedure in **PASS**.

Power and Sample Size of the Bridging Study

Chow, Shao, and Hu (2002) define $\hat{P}(\Delta T(x))$ as the probability of generalizability of the results of the original population to the new population.

If it is determined that a bridging study must be conducted, $\hat{P}(\Delta T^*(x))$ estimates the power of that study at a given sample size. Using a simple search algorithm, this power estimate can be used to determine an appropriate sample size for the bridging study.

Example 1 – Finding Sample Size

Chow, Shao, and Hu (2002) page 394 present the following example. A certain drug has been cleared for use in the originally studied region using a two-sample t-test from a design with both a treatment group and a control group. The trial resulted in the following summary statistics:

$N_{OT} = 30$	$\bar{x}_{OT} = 19.8$	$s_{OT} = 8.4$
$N_{OC} = 30$	$\bar{x}_{OC} = 26.5$	$s_{OC} = 9.3$

Sponsors in another region not included in the original study would like to register this drug for use in that region as well. They want to assess the generalizability if the sensitivity index, Δ , is between 0.7 and 0.9. They also want to determine the reduction probability of the original study, so they add $\Delta = 1.0$ to the list.

In this new region, the policy is that a bridging study is not necessary if there is no more than a 10% reduction in the reproducibility probability of the original study.

If a bridging study is found necessary, they would like to determine sample necessary to obtain a power of 0.7 with a significance level of 0.05. They are planning a balanced study.

Setup

If the procedure window is not already open, use the PASS Home window to open it. The parameters for this example are listed below and are stored in the **Example 1** settings file. To load these settings to the procedure window, click **Open Example Settings File** in the Help Center or File menu.

Design Tab

Solve For	. Sample Size
Noт (Sample Size of Group от)	.30
Noc (Sample Size of Group oc)	.30
Do (Mean Difference)	6.7
Sot (Std Deviation of Group ot)	.8.4
Soc (Std Deviation of Group oc)	.9.3
Power of Bridging Study	.0.7
Alpha	.0.05
Group Allocation	. Equal (Nвт = Nвс)
Δ (Sensitivity Index)	.0.7 0.75 0.8 0.85 0.9 1

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Reports

Numeric Res	sults	
Solve For: Hypotheses:	Sample Size H0: µBT - µBC = 0 vs.	Н1: µвт - µвс ≠ 0

						Original Study							
Sample Size Sensitivity								nple	Mean	Standard Deviation			
Power		Nвс	NB	Index	Alpha	Power'	 Nот	Noc	Difference	Treatment Sот	Control Soc		
0.70923	46	46	92	0.70	0.05	0.52230	30	30	-6.7	8.4	9.3		
0.70702	40	40	80	0.75	0.05	0.57920	30	30	-6.7	8.4	9.3		
0.70351	35	35	70	0.80	0.05	0.63450	30	30	-6.7	8.4	9.3		
0.70180	31	31	62	0.85	0.05	0.68713	30	30	-6.7	8.4	9.3		
0.70563	28	28	56	0.90	0.05	0.73620	30	30	-6.7	8.4	9.3		
0.70807	23	23	46	1.00	0.05	0.82109	30	30	-6.7	8.4	9.3		

Power The probability of rejecting a false null hypothesis when the alternative hypothesis is true.

NBT The number of bridging study subjects assigned to the treatment group.

NBC The number of bridging study subjects assigned to the control group.

NB The total sample size of the bridging study.

 Δ The sensitivity index. This is the ratio of the signal-to-noise ratios in the bridging study and the original study.

Alpha The probability of rejecting a true null hypothesis.

Power' The power of the test in the original study assuming that the parameter values are the sample values and that the noncentrality parameter is $\Delta \times T$. This value is used to determine whether the bridging study is necessary.

Not The number of subjects assigned to the treatment group in the original study.

Noc The number of subjects assigned to the control group in the original study.

Diff The difference between the group means $(\tau - c)$ in the original study.

Som The response standard deviation of subjects assigned to the treatment group in the original study.

Soc The response standard deviation of subjects assigned to the control group in the original study.

Summary Statements

A parallel two-group bridging study design will be used in an attempt reproduce the significant mean difference result of the original study in the bridging study region. The bridging study comparison will be made using a two-sided t-test, with a Type I error rate (α) of 0.05. The group sample sizes of the original study were 30 (treatment) and 30 (control). The within-group standard deviations of the original study were 8.4 (treatment) and 9.3 (control). The original study mean difference (treatment minus control) was -6.7. For a sensitivity index (ratio of the bridging study to original study signal-to-noise ratios) of 0.7 with reproducibility probability (power) 70%, the number of subjects needed for the bridging study will be 46 in the treatment group and 46 in the control group.

Dropout-Inflated Sample Size

Dropout Rate	Sample Size			Dro E S	pout-Infla Enrollmen ample Siz	Expected Number of Dropouts			
	Nвт	Nвс	Νв	Nвт'	Nвс'	NB'	Dτ	Dc	D
20%	46	46	92	58	58	116	12	12	24
20%	40	40	80	50	50	100	10	10	20
20%	35	35	70	44	44	88	9	9	18
20%	31	31	62	39	39	78	8	8	16
20%	28	28	56	35	35	70	7	7	14
20%	23	23	46	29	29	58	6	6	12
Dropout Rate	The percen study and	tage of sul I for whom	ojects (or ite no respons	ems) that are o e data will be	expected to collected (i	be lost at ra .e., will be tre	ndom durin eated as "m	g the cou nissing").	irse of Abbre

will
Nвт vays
Nв vay

Dropout Summary Statements

Anticipating a 20% dropout rate, 58 subjects should be enrolled in Group 1, and 58 in Group 2, to obtain final group sample sizes of 46 and 46, respectively.

References

Shao, J. and Chow, S.C. 2002. 'Reproducibility probability in clinical trials.' Statistics in Medicine, Volume 21, Pages 1727-1742.

Chow, S.C., Shao, J. and Hu, O.Y.P. 2002. 'Assessing sensitivity and similarity in bridging studies.' Journal of Biopharmaceutical Statistics. Volume 12(3), Pages 385-400.

Chow, S.C., Shao, J., Wang, H., and Lokhnygina, Y. 2018. Sample Size Calculations in Clinical Research, Third Edition. Taylor & Francis/CRC. Boca Raton, Florida.

This report shows the sample size for the indicated parameter configurations.

The first question to be answered is whether a bridging study is necessary. The last row of the report shows the reproducibility probability (Power') to be 0.82109. The policy in the new region is that Power' be no less than 10% below this value. This results in a cutoff of 0.82109 \times 0.9 = 0.73898. The Power' for Δ = 0.9 is 0.73620 which is still below the cutoff of 0.73898. Hence, for any of the sensitivity index values between 0.7 and 0.9, the decision would be to run the bridging study. The N_B column gives the required sample size to achieve 70% power. For example, if the anticipated value of Δ = 0.8, a sample size of 35 per group is required.

Plots Section



This plot shows the sample size versus sensitivity index.

Example 2 – Validation using Chow, Shao, and Hu (2002)

Chow, Shao, and Hu (2002) page 394 present the following example which we will use to validate this procedure. A certain drug has been cleared for use in the originally studied region using a two-sample t-test from a design with both a treatment group and a control group. The trial resulted in the following summary statistics as shown in Table 4:

$N_{OT} = 30$	$\bar{x}_{OT} = 19.8$	$s_{OT} = 8.4$
$N_{OC} = 30$	$\bar{x}_{OC} = 26.5$	$s_{OC} = 9.3$

Sponsors in another region not included in the original study would like to register this drug for use in that region as well. They want to assess the generalizability if the sensitivity index, Δ , is between 0.7 and 0.85.

In Table 5 on page 394 obtained the following sample sizes for Δ between 0.7 and 0.85: 92, 80, 70, and 62.

Setup

If the procedure window is not already open, use the PASS Home window to open it. The parameters for this example are listed below and are stored in the **Example 2** settings file. To load these settings to the procedure window, click **Open Example Settings File** in the Help Center or File menu.

Design Tab	
Solve For	Sample Size
Noт (Sample Size of Group от)	30
Noc (Sample Size of Group oc)	
Do (Mean Difference)	6.7
Sot (Std Deviation of Group ot)	8.4
Soc (Std Deviation of Group oc)	9.3
Power of Bridging Study	0.7
Alpha	0.05
Group Allocation	Equal (Nвт = Nвс)
Δ (Sensitivity Index)	0.7 0.75 0.8 0.85

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results

```
Solve For:
Sample Size

Hypotheses:
H0: μвт - μвс = 0
vs.
H1: μвт - μвс ≠ 0
```

		Duidai					Original Study					
Sample Size Sensitivity							Sample		Mean	Standard Deviation		
Power		Nвс	Νв	Index Δ	Alpha	Power'	 Nот	Noc	Difference	Treatment Sот	Control Soc	
0.70923	46	46	92	0.70	0.05	0.52230	30	30	-6.7	8.4	9.3	
0.70702	40	40	80	0.75	0.05	0.57920	30	30	-6.7	8.4	9.3	
0.70351	35	35	70	0.80	0.05	0.63450	30	30	-6.7	8.4	9.3	
0.70180	31	31	62	0.85	0.05	0.68713	30	30	-6.7	8.4	9.3	

Plots



PASS has also calculated sample sizes of 92, 80, 70, and 62 so the procedure is validated. The plot shows the sample size versus sensitivity index.