

Chapter 259

Bridging Study using a Non-Inferiority Test of Two Groups (Continuous Outcome)

Introduction

This procedure calculates the power and sample size required for bridging studies that use a non-inferiority test of the mean difference between the results for the two regions: original and bridging. Only a brief introduction to the subject will be given here. For a comprehensive discussion, refer to Liu, Hsueh, and Chen (2002).

Bridging Studies

Once a pharmaceutical product has been approved for use in one or more regions (countries) through a set of clinical trials, it is often desirable to register the product in a new region that was not included in the original study. When the cost and time needed to complete an additional set of clinical trials in the new region is prohibitive, a *bridging methodology* may be used to obtain the approval. The bridging analysis compares the results of a smaller and shorter *bridging study* in the new region with the data obtained in the original study.

The bridging analysis makes use of a two-group design in which the effectiveness in the new region is compared to the effectiveness in the original region using a non-inferiority test. The effectiveness in each region is measured by the difference between the means of a treatment group and a control group. The non-inferiority test shows that the differences in the two regions do not differ by more than a small amount, called the non-inferiority.

Test Statistics

This section summarizes the results found in Liu, Hsueh, and Chen (2002), page 974 - 976.

Original Study

Let Y_{ijk} be the clinical response of subject k on receiving treatment j in original study i . It is assumed that $i = 1, \dots, I$. Also, $j = T$ (treatment), C (control) and $k = 1, \dots, N_{ij}$. Hence Y_{ijk} includes the response data from each of the original trials. Assume that the Y_{ijk} are independently normally distributed with means μ_{ij} and variance σ_{ij}^2 . Further assume that μ_{ij} has a normal distribution with mean μ_{0j} and variance γ_{0j}^2 . Hence, the Y_{ijk} 's are independently normally distributed with mean μ_{0j} and variance $\omega_{ij}^2 = \sigma_{ij}^2 + \gamma_{0j}^2$.

Let \bar{Y}_{ij} be the sample means. The MLE of μ_{0j} is

$$t_{0j} = \frac{\sum Y_{ij} / (w_{ij}^2 / N_{ij})}{\sum 1 / (w_{ij}^2 / N_{ij})}, i = 1, \dots, I; j = T, C$$

Bridging Study using a Non-Inferiority Test of Two Groups (Continuous Outcome)

where

$$w_{ij}^2 = \sum \frac{(Y_{ijk} - t_{0j})^2}{N_{ij}}$$

is the MLE of ω_{ij}^2 . The MLE's t_{0j} and ω_{ij}^2 are solved for iteratively.

Bridging Study

Let Y_{Bjk} be the clinical response of subject k on receiving treatment j in the bridging study conducted in the new region. It is assumed that $j = T, C$ and $k = 1, \dots, N_{Bj}$. As before, the Y_{Bjk} 's are independently normally distributed with mean μ_{Bj} and variance ω_{Bj}^2 .

The MLE of μ_{Bj} is the sample mean Y_{Bj} . Let $t_{Bj} = Y_{Bj}, j = T, C$.

Non-Inferiority Test

The MLEs t_{0j} and t_{Bj} are independently normally distributed with asymptotic variances estimated by

$$s_{0j}^2 = \frac{1}{\sum 1/(w_{ij}^2/N_{ij})}$$

and

$$s_{Bj}^2 = \sum \frac{(Y_{Bjk} - t_{Bj})^2}{N_{Bj}^2}$$

Let $E_L = -NIM$ be the lower non-inferiority limit for the mean difference between regions, assuming $NIM > 0$. NIM is the non-inferiority margin. Often, NIM is set using $NIM = f(t_{0T} - t_{0C})$ where f is between 0 and 0.5.

The non-inferiority hypotheses, assuming higher values are better, are

$$H_0: \theta \leq -NIM \text{ vs } H_1: \theta > -NIM$$

where

$$\theta = (\mu_{BT} - \mu_{BC}) - (\mu_{OT} - \mu_{OC})$$

is the difference in treatment effects between the two regions.

The test statistic

$$t = (t_{BT} - t_{BC}) - (t_{OT} - t_{OC})$$

is an asymptotically unbiased estimate for θ .

The variance of t is given by

$$s^2 = s_{BT}^2 + s_{BC}^2 + s_{OT}^2 + s_{OC}^2.$$

Bridging Study using a Non-Inferiority Test of Two Groups (Continuous Outcome)

The test statistic for the non-inferiority test is

$$T_L = \frac{(t + NIM)}{s}$$

The null hypothesis is rejected, and non-inferiority is concluded at significance level α if and only if $T_L > z_\alpha$, where z_α is the α^{th} upper percentile of the standard normal distribution. For a one-sided test such as this, α is often set to 0.025.

Power Calculation

Based on the above results, Liu *et al.* (2002) estimate the sample size required to meet the power, significance level, and effect size requirement as

$$N_B \geq \frac{A_1}{A_2 - A_3}$$

where

$$A_1 = \frac{\sigma_{BT}^2}{g_{BT}} + \frac{\sigma_{BC}^2}{1 - g_{BT}}$$

$$A_2 = \frac{NIM^2}{(z_\alpha + z_\beta)^2}$$

$$A_3 = s_{OT}^2 + s_{OC}^2$$

$$g_{BT} = \frac{N_{BT}}{N_B}$$

where $\beta = 1 - \text{Power}$, σ_{BT}^2 is often estimated by s_{OT}^2 , σ_{BC}^2 is often estimated by s_{OC}^2 , and the actual difference between the two study differences is zero.

The power is obtained by rearranging this formula.

Example 1 – Finding Sample Size

A certain drug has been cleared for use in North America using parallel-group, treatment versus control clinical trials. These trials resulted in the following summary statistics:

$$N_{OT} = 973 \quad \hat{\mu}_{OT} = 15.47 \quad s_{OT} = 11.86$$

$$N_{OC} = 948 \quad \hat{\mu}_{OC} = 4.14 \quad s_{OC} = 10.39$$

Researchers in a region not included in the original study would like to register the new drug for use in that region. To do so, they are planning a bridging study with a significance level of 0.05 and a power of 0.8. They will set $\sigma_{BT} = s_{OT}$ and $\sigma_{BC} = s_{OC}$. They want to calculate the necessary sample size when f is 0.2, 0.3, or 0.4. They are planning a balanced study.

Setup

If the procedure window is not already open, use the PASS Home window to open it. The parameters for this example are listed below and are stored in the **Example 1** settings file. To load these settings to the procedure window, click **Open Example Settings File** in the Help Center or File menu.

Design Tab

Solve For	Sample Size
N _{OT} (Sample Size of Group OT).....	973
N _{OC} (Sample Size of Group OC)	948
Do (Mean Difference).....	11.33
S _{OT} (Std Deviation of Group OT).....	11.86
S _{OC} (Std Deviation of Group OC)	10.39
Power.....	0.8
Alpha.....	0.025
Group Allocation	Equal (N_{BT} = N_{BC})
Non-Inferiority Margin Input	Enter f, the proportion NIM is of Do
f (Proportion NIM is of Do)	0.2 0.3 0.4
σ _{BT} (Std Deviation of Group BT).....	11.86
σ _{BC} (Std Deviation of Group BC).....	10.39

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Reports

Numeric Results

Solve For: [Sample Size](#)
 Definition: $\theta = (\mu_{BT} - \mu_{BC}) - (\mu_{OT} - \mu_{OC})$
 Hypotheses: $H_0: \theta \leq -NIM$ vs. $H_1: \theta > -NIM$
 H1 Assumption: $\theta = 0$

Bridging Study									Original Study				
Power	N _{BT}	N _{BC}	N _B	Non-Inferiority		Std Dev		Alpha	N _{OT}	N _{OC}	Mean Diff Do	Std Dev	
				Prop f	Value NIM	Trt σ_{BT}	Ctrl σ_{BC}					Trt S _{OT}	Ctrl S _{OC}
0.80031	629	629	1258	0.2	2.266	11.86	10.39	0.025	973	948	11.33	11.86	10.39
0.80021	205	205	410	0.3	3.399	11.86	10.39	0.025	973	948	11.33	11.86	10.39
0.80195	106	106	212	0.4	4.532	11.86	10.39	0.025	973	948	11.33	11.86	10.39

Power The probability of rejecting a false null hypothesis when the alternative hypothesis is true.
 N_{BT} The number of bridging study subjects assigned to the treatment group.
 N_{BC} The number of bridging study subjects assigned to the control group.
 N_B The total sample size of the bridging study.
 f The proportion of |Do| used as the magnitude of the non-inferiority margin. $NIM = f \times |Do|$.
 NIM The magnitude of the non-inferiority margin. $L = -NIM$ or $U = NIM$. $NIM > 0$.
 σ_{BT} The response standard deviation of subjects assigned to the treatment group in the bridging study.
 σ_{BC} The response standard deviation of subjects assigned to the control group in the bridging study.
 Alpha The probability of rejecting a true null hypothesis.
 N_{OT} The number of subjects assigned to the treatment group in the original study.
 N_{OC} The number of subjects assigned to the control group in the original study.
 Do The difference between the group means ($\tau - c$) in the original study.
 S_{OT} The response standard deviation of subjects assigned to the treatment group in the original study.
 S_{OC} The response standard deviation of subjects assigned to the control group in the original study.

Summary Statements

The bridging study sample sizes of 629 in the treatment group and 629 in the control group achieve 80% power using a non-inferiority test of the difference between the two group means. The significance level (alpha) of the non-inferiority test is 0.025. The magnitude of the non-inferiority margin is 2.266. The bridging-study treatment-group standard deviation is 11.86. The bridging-study control-group standard deviation is 10.39. The summary statistics of the original study are as follows. The treatment group sample size was 973. The control group sample size was 948. The difference between the group means was 11.33. The treatment group standard deviation was 11.86. The control group standard deviation was 10.39.

Bridging Study using a Non-Inferiority Test of Two Groups (Continuous Outcome)

Dropout-Inflated Sample Size

Dropout Rate	Sample Size			Dropout-Inflated Enrollment Sample Size			Expected Number of Dropouts		
	N _{BT}	N _{BC}	N _B	N _{BT'}	N _{BC'}	N _{B'}	D _T	D _C	D
20%	629	629	1258	787	787	1574	158	158	316
20%	205	205	410	257	257	514	52	52	104
20%	106	106	212	133	133	266	27	27	54

- Dropout Rate The percentage of subjects (or items) that are expected to be lost at random during the course of the study and for whom no response data will be collected (i.e., will be treated as "missing"). Abbreviated as DR.
- N_{BT}, N_{BC}, and N_B The evaluable sample sizes at which power is computed (as entered by the user). If N_{BT} and N_{BC} subjects are evaluated out of the N_{BT'} and N_{BC'} subjects that are enrolled in the study, the design will achieve the stated power.
- N_{BT'}, N_{BC'}, and N_{B'} The number of subjects that should be enrolled in the study in order to obtain N_{BT}, N_{BC}, and N_B evaluable subjects, based on the assumed dropout rate. N_{BT'} and N_{BC'} are calculated by inflating N_{BT} and N_{BC} using the formulas $N_{BT'} = N_{BT} / (1 - DR)$ and $N_{BC'} = N_{BC} / (1 - DR)$, with N_{BT'} and N_{BC'} always rounded up. (See Julious, S.A. (2010) pages 52-53, or Chow, S.C., Shao, J., Wang, H., and Lohknygina, Y. (2018) pages 32-33.)
- D_T, D_C, and D The expected number of dropouts. $D_T = N_{BT'} - N_{BT}$, $D_C = N_{BC'} - N_{BC}$, and $D = D_T + D_C$.

Dropout Summary Statements


Anticipating a 20% dropout rate, 787 subjects should be enrolled in Group 1, and 787 in Group 2, to obtain final group sample sizes of 629 and 629, respectively.

References

Liu, J.P., Hsueh, H., Chen, J.J. 2002. 'Sample Size Requirements for Evaluation of Bridging Evidence.' Biometrical Journal, Volume 44 (8), Pages 969-981.

This report shows the sample size for the indicated parameter configurations.

Plots Section



This plot shows the power versus the sample size. Note that we had to reduce the font size of the subtitle so that it would fit in the space allotted.

Example 2 – Validation using Liu et al. (2002)

Liu et al. (2002) include Table 2 of example results on page 977. We will use the entry from the sixth row and second column ($f = 0.2$) of this table as our validation example. The other table parameters are $CV = 80\%$, $N_O = 1000$, $g_{NT} = 0.5$. They find the resulting bridging study sample size to be 55 per group (110 total).

These input values are consistent with the following summary statistics:

$$N_{OT} = 500 \quad \hat{\mu}_{OT} = 4 \quad s_{OT} = 0.8$$

$$N_{OC} = 500 \quad \hat{\mu}_{OC} = 2 \quad s_{OC} = 0.8$$

The significance level = 0.05 and the power = 0.8. Set $\sigma_{BT} = s_{OT}$ and $\sigma_{BC} = s_{OC}$.

Setup

If the procedure window is not already open, use the PASS Home window to open it. The parameters for this example are listed below and are stored in the **Example 2** settings file. To load these settings to the procedure window, click **Open Example Settings File** in the Help Center or File menu.

Design Tab

Solve For	Sample Size
N _{OT} (Sample Size of Group o _T).....	500
N _{OC} (Sample Size of Group o _C)	500
Do (Mean Difference).....	2
S _{OT} (Std Deviation of Group o _T).....	0.8
S _{OC} (Std Deviation of Group o _C)	0.8
Power.....	0.8
Alpha.....	0.05
Group Allocation	Equal (N_{BT} = N_{BC})
Non-Inferiority Margin Input	Enter f, the proportion NIM is of Do
f (Proportion NIM is of Do)	0.2
σ _{BT} (Std Deviation of Group b _T).....	0.8
σ _{BC} (Std Deviation of Group b _C).....	0.8

Bridging Study using a Non-Inferiority Test of Two Groups (Continuous Outcome)

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results

Solve For: [Sample Size](#)
 Definition: $\theta = (\mu_{BT} - \mu_{BC}) - (\mu_{OT} - \mu_{OC})$
 Hypotheses: $H_0: \theta \leq -NIM$ vs. $H_1: \theta > -NIM$
 H1 Assumption: $\theta = 0$

Bridging Study									Original Study				
Power	NBT	NBC	NB	Non-Inferiority		Std Dev		Alpha	Not	Noc	Mean Diff Do	Std Dev	
				Prop f	Value NIM	Trt σ_{BT}	Ctrl σ_{BC}					Trt σ_{OT}	Ctrl σ_{OC}
0.80063	55	55	110	0.2	0.4	0.8	0.8	0.05	500	500	2	0.8	0.8

PASS has also calculated a sample size of 55 per group, so the procedure is validated.