

Chapter 364

Mixed Models Tests for Slope-Interaction in a 2x2 Factorial 3-Level Hierarchical Design with Random Slopes (Level-2 Randomization)

Introduction

This procedure calculates power and sample size for a three-level hierarchical mixed model which is randomized at the **second** level. The associated **longitudinal** study uses a 2-by-2 factorial design with two binary factors X and Z, each with two possible values (0 and 1). This results in four treatment arms. The goal of the study is to test whether the slopes of subjects across time are different from what would be expected if the effect of the two factors were additive. That is, one wants to test the three-way interaction between the two binary factors and time.

In most cases, this design is called a *repeated measures* design. The classic example is a study in which the level-2 units are subjects which are nested in level-3 units (e.g., classes, clinics, or hospitals). The level-1 units are time points at which measurements are taken. This factor is nested in the level-2 units.

This procedure is for longitudinal studies in which each subject is measured two or more times.

In this case of level-2 randomization, each level-2 unit (subject) is randomly assigned to one of the four treatments combinations. Hence, a level-3 unit will include subjects in all four treatment groups.

Each subject is assumed to have a separate, *random* slope.

Technical Details

Our formulation comes from Ahn, Heo, and Zhang (2015), chapter 6, section 6.6.2, pages 221-222. The hierarchical mixed model that is adopted is

$$Y_{ijk} = \beta_0 + \delta_X X_{ijk} + \delta_Z Z_{ijk} + \delta_T T_{ijk} + \delta_{XZ} X_{ijk} Z_{ijk} + \delta_{XT} X_{ijk} T_{ijk} + \delta_{ZT} Z_{ijk} T_{ijk} + \delta_{XZT} X_{ijk} Z_{ijk} T_{ijk} + v_{j(i)} T_{ijk} + u_i + u_{j(i)} + e_{ijk}$$

where

- Y_{ijk} is the continuous response of the k^{th} level-1 unit, within the j^{th} level-2 unit, within the i^{th} level-3 unit.
- X_{ijk} is an indicator variable that is equal to “1” if the j^{th} level-2 unit is assigned to receive intervention X and “0” otherwise. Thus, $X_{ijk} = X_j$ for all i and k .
- Z_{ijk} is an indicator variable that is equal to “1” if the j^{th} level-2 unit is assigned to receive intervention Z and “0” otherwise. Thus, $Z_{ijk} = Z_j$ for all i and k .
- β_0 is the fixed intercept.
- δ_X is the treatment effect of factor X.
- δ_Z is the treatment effect of factor Z.
- δ_{XZ} is the interaction effect of factors X and Z.
- δ_{XT} is the interaction effect of factors X and Z.
- δ_{ZT} is the interaction effect of factors X and Z.
- δ_{XZT} is the 3-way interaction effect of X, Z, and time. **The is the coefficient of interest.**
- u_i is the level-3 random intercept effect for the i^{th} level-3 unit. It is distributed as $N(0, \sigma_3^2)$.
- $u_{j(i)}$ is the level-2 random intercept effect for the $j(i)^{\text{th}}$ level-2 unit. It is distributed as $N(0, \sigma_2^2)$.
- e_{ijk} is the level-1 random intercept effect that is distributed as $N(0, \sigma_e^2)$.
- σ_e^2 is variance of the subject-specific random slopes.
- σ^2 is the variance of Y when slopes are fixed, where $\sigma^2 = \sigma_3^2 + \sigma_2^2 + \sigma_e^2$.
- ρ_1 is the correlation among level-1 units which are in a particular level-2 unit.
- ρ_2 is the correlation among level-2 units which are in a particular level-3 unit.
- $K_{0,0}$ is the number of level-2 units per level-3 unit for which $X = 0$ and $Z = 0$.
- $K_{0,1}$ is the number of level-2 units per level-3 unit for which $X = 0$ and $Z = 1$.
- $K_{1,0}$ is the number of level-2 units per level-3 unit for which $X = 1$ and $Z = 0$.
- $K_{1,1}$ is the number of level-2 units per level-3 unit for which $X = 1$ and $Z = 1$.
- C is the number of level-3 units.
- M is the number of level-1 units per level-2 unit. It is the number of measurement times.

The test of significance of the product $X_{ijk} Z_{ijk} T_{ijk}$ is the interaction effect of X, Z, and Time.

This is the test statistic of interest. It tests whether the subject-specific slopes behave the same across all treatment combinations.

Assume that δ_{XZT} is to be tested using a Wald test. The statistical hypotheses are $H_0: \delta_{XZT} = 0$ vs. $H_a: \delta_{XZT} \neq 0$.

Mixed Models Tests for Slope-Int'n in a 2x2 Fact. 3-Lvl Hier. Design with Rand. Slopes (Lvl-2 Rand.)

The power is calculated using

$$Power = \Phi \left\{ \left| \frac{\delta_{XZT}}{\sigma} \right| \sqrt{\frac{CK_{0,0}M \text{Var}(T)}{f \left(\frac{1}{K_{0,0}} + \frac{1}{K_{1,1}} + \frac{1}{K_{1,0}} + \frac{1}{K_{0,1}} \right)}} - \Phi(1 - \alpha/2) \right\}$$

where $f = 1 - \rho_1 + r\tau M \text{Var}(T)$ and $\text{Var}(T) = \sum_{j=1}^M (T_j - \bar{T})^2 / M$.

Procedure Options

This section describes the options that are specific to this procedure. These are located on the Design tab. For more information about the options of other tabs, go to the Procedure Window chapter.

Design Tab

The Design tab contains most of the parameters and options that you will be concerned with.

Solve For

Solve For

This option specifies the parameter to be solved for from the other parameters. The parameters that may be selected are δ_{XZT} (Three-Way Interaction), Power, C, K00, or M.

Note that the value selected here always appears as the vertical axis on the charts.

The program is set up to calculate power directly. To find appropriate values of the other parameters, a binary search is made using an iterative procedure until an appropriate value is found. This search considers integer values of K00, C, and M only.

Power and Alpha

Power

This option specifies one or more values for power. Power is the probability of rejecting a false null hypothesis, and is equal to one minus Beta. Beta is the probability of a type-II error, which occurs when a false null hypothesis is not rejected.

Values must be between zero and one. Historically, the value of 0.80 (Beta = 0.20) was used for power. Now, 0.90 (Beta = 0.10) is also commonly used.

A single value may be entered or a range of values such as *0.8 to 0.95 by 0.05* may be entered.

If your only interest is in determining the appropriate sample size for a confidence interval, set power to 0.5.

Alpha

This option specifies one or more values for the probability of a type-I error. A type-I error occurs when a true null hypothesis is rejected.

Values must be between zero and one. Usually, the value of 0.05 is used for alpha and this has become a standard. This means that about one test in twenty will falsely reject the null hypothesis. You should pick a value for alpha that represents the risk of a type-I error you are willing to take in your experimental situation.

You may enter a range of values such as *0.01 0.05 0.10* or *0.01 to 0.10 by 0.01*.

Sample Size – Number of Level-1, Level-2, and Level-3 Units

C (Level-3 Units)

This is the number of level-2 units (usually subjects or patients) per level-3 unit. For example, if the three levels are repeated measures in subjects in doctors, this level is the number of subjects per doctor.

This value must be a positive number that is at least 1.

You can use a list of values such as "10 15 20". A separate analysis will be run for each element in the list.

K00 (Group 00 Count (X=0, Z=0))

This is the number of level-2 units per level-3 unit in group 00, the (0,0) cell in a 2-by-2 factorial table of X and Z. The total sample size (number of level-1 units) of this group is $K00 \times C \times M$.

Note that this is a longitudinal study.

Each of the factors X and Z have two levels: 0 (no intervention) and 1 (intervention). Thus, in this group in which $X = 0$ and $Z = 0$, neither of the interventions occur. Hence, this is called the control group.

This value must be a positive integer.

You can enter a list of values such as "10 20 30". A separate analysis will be run for each element in the list.

K01 (Group 01 Count (X=0, Z=1))

This is the number of level-2 units per level-3 unit in group 01, the (0,1) cell in a 2-by-2 factorial table of X and Z. The total sample size (number of level-1 units) of this group is equal to $K01 \times C \times M$.

Note that this is a longitudinal study.

Each of the factors X and Z have two levels: 0 (no intervention) and 1 (intervention). Thus, in this group in which $X = 0$ and $Z = 1$, an intervention occurs for the second factor but not the first.

Using Multiples of K00

If you simply want a multiple of the value for group 00, enter the multiple followed by "K00", with no blanks. If you want to use K00 directly, you do not have to enter a leading "1".

For example, all of the following are valid entries: 10K00 2K00 0.5K00 K00.

You can use a list of values such as "10 20 30" or "K00 2K00 3K00".

K10 (Group 01 Count (X=1, Z=0))

This is the number of level-2 units per level-3 unit in group 10, the (1,0) cell in a 2-by-2 factorial table of X and Z. The total sample size (number of level-1 units) of this group is equal to $K10 \times C \times M$.

Note that this is a longitudinal study.

Each of the factors X and Z have two levels: 0 (no intervention) and 1 (intervention). Thus, in this group in which $X = 1$ and $Z = 0$, an intervention occurs for the first factor but not the second.

Mixed Models Tests for Slope-Int'n in a 2x2 Fact. 3-Lvl Hier. Design with Rand. Slopes (Lvl-2 Rand.)**Using Multiples of K00**

If you simply want a multiple of the value for group 00, enter the multiple followed by "K00", with no blanks. If you want to use K00 directly, you do not have to enter a leading "1".

For example, all of the following are valid entries: 10K00 2K00 0.5K00 K00.

You can use a list of values such as "10 20 30" or "K00 2K00 3K00".

K11 (Group 11 Count (X=1, Z=1))

This is the number of level-2 units per level-3 unit in group 11, the (1,1) cell in a 2-by-2 factorial table of X and Z. The total sample size (number of level-1 units) of this group is equal to $K11 \times C \times M$.

Note that this is a longitudinal study.

Each of the factors X and Z have two levels: 0 (no intervention) and 1 (intervention). Thus, in this group in which $X = 1$ and $Z = 1$, both interventions occur.

Using Multiples of K00

If you simply want a multiple of the value for group 00, enter the multiple followed by "K00", with no blanks. If you want to use K00 directly, you do not have to enter a leading "1".

For example, all of the following are valid entries: 10K00 2K00 0.5K00 K00.

You can use a list of values such as "10 20 30" or "K00 2K00 3K00".

M (Level-1 Unit Count Per Level-2 Unit)

This is the number of level-1 units (e.g., time points) per level-2 unit (e.g., subject) in all four groups.

This value must be a positive number that is at least 2.

List

You can use a list of values such as "3 5 6". A separate analysis will be run for each element in the list.

Effect Size **δ_{XZT} (Three-Way Interaction)**

Enter one or more values for δ_{XZT} , the regression coefficient of the $X_{ijk} \times Z_{ijk} \times T_{ijk}$ term in the mixed model. It is this interaction among the four group slopes at which the study is to be powered. That is, the power is the probability of detecting an interaction of at least this amount. This value is not the true interaction. Rather, it is the interaction that you want to be able to detect.

Slope Interaction

Suppose we define β_{xz} to be the slope of the level-1 units in the (x,z) cell in a 2-by-2 factorial table. For example, β_{10} is the slope of the data in the (1,0) cell in a 2-by-2 factorial table of X and Z.

This interaction is a difference of slope differences. It is constructed as the difference between the two slope differences ($\beta_{11} - \beta_{10}$) and ($\beta_{01} - \beta_{00}$). It is the failure of the two interventions to act independently of each other.

δ_{XZT} can be any non-zero value (positive or negative). Since this procedure uses a two-sided test statistic, you will get the same result with either positive or negative values.

Mixed Models Tests for Slope-Int'n in a 2x2 Fact. 3-Lvl Hier. Design with Rand. Slopes (Lvl-2 Rand.)**Syntax**

You can enter a single value such as 10 or a series of values such as 10 20 30 or 5 to 50 by 5. When a series of values is entered, PASS will generate a separate calculation result for each value of the series.

 σ (Standard Deviation)

Enter the standard deviation of Y_{ijk} assuming the slopes are fixed. If the slopes are assumed random, you need to obtain the estimated variance from the fit of a fixed-slope model. You might obtain variance component estimates for the quantities used to form σ^2 from previous studies or a pilot study.

Definition of σ^2

For clarity, in the following we are using square-brackets to enclose the subscripts.

Let

$V(e)$ = Variance of error term, $e[ijk]$.

$V(2)$ = Variance of level-2 random intercept, $u[j(i)]$.

$V(3)$ = Variance of level-3 random intercept, $u[i]$.

$V(\tau)$ = Variance of random slopes. If the slopes are fixed, $V(\tau) = 0$.

$T[k]^2 = T[k] \times T[k]$.

σ^2 is defined as $\sigma^2 = V(3) + V(2) + V(e)$.

Note that for random-slope models $V(Y[ijk]) = \sigma^2 + V(\tau) \times T[k]^2$ so that in the random slope case, you cannot simply use $V(Y[ijk])$ for σ^2 .

Range

σ must be a positive number.

Syntax

You can enter a single value such as 5 or a series of values such as 1 3 5 7 9 or 1 to 10 by 1. When a series of values is entered, PASS will generate a separate calculation result for each value of the series.

Help for Estimating σ

Press the small ' σ ' button to the right to obtain calculation options for estimating the standard deviation.

 ρ_1 (Correlation Among Level 1 Units)

This is the correlation of each pair of level 1 units in a particular level 2 unit.

Possible values are from 0 to just below 1. Typical values are between 0.0001 and 0.5. You may enter a single value or a list of values.

 $r\tau (V(\tau)/\sigma^2)$

Enter the ratio of the variance of subject-specific random slopes, $V(\tau)$, to the sum of the other variances, σ^2 . You must estimate this value using data from previous studies or a pilot study.

Possible values are from 0 to just below 1. You may enter a single value or a list of values.

Example 1 – Calculating Power

Suppose that a three-level hierarchical design is planned in which there will be two interventions. Each intervention will be whether one of two drugs is administered. This will result in four treatment groups. There will be five measurements per subject and the four treatments will be randomly applied to level-2 units.

The analysis will be a mixed model of continuous data using the model given earlier in this chapter. The subject-specific slopes will be assumed to be random. The goal of the study is to test the regression coefficient of the XZT interaction. The following parameter settings are to be used for the power analysis: $\delta_{XZT} = 3$; $\sigma = 9.8$; $\rho_1 = 0.1$; $\tau = 0.1$; $C = 5$ or 10 ; $M = 5$; $\alpha = 0.05$; and $K_{00} = K_{01} = K_{10} = K_{11} = 5$ to 20 by 5 . Find the power of each combination of parameter settings.

Setup

This section presents the values of each of the parameters needed to run this example. First, from the PASS Home window, load the **Mixed Models Tests for Slope-Int'n in a 2x2 Fact. 3-Lvl Hier. Design with Rand. Slopes (Lvl-2 Rand.)** procedure window. You may then make the appropriate entries as listed below, or open **Example 1** by going to the **File** menu and choosing **Open Example Template**.

<u>Option</u>	<u>Value</u>
Design Tab	
Solve For	Power
Alpha.....	0.05
C (Level 3 Units)	5 10
K00 (Group 00 Count (X=0, Z=0))	5 10 15 20
K01 (Group 01 Count (X=0, Z=1))	K00
K10 (Group 10 Count (X=1, Z=0))	K00
K11 (Group 11 Count (X=1, Z=1))	K00
M (Level 1 Units Per Level-2 Unit).....	5
δ_{XZT} (Three-Way Interaction)	3
σ (Standard Deviation)	9.8
ρ_1 (Correlation Among Level-1 Units)	0.1
τ ($V(\tau) / \sigma^2$)	0.1

Annotated Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results

Numeric Results for an Interaction in a 2x2 Factorial												
	Total		Grp00	Grp01	Grp10	Grp11		2 x 2		ICC		
	Lvl 1	Lvl 3	Lvl 2	Lvl 2	Lvl 2	Lvl 2	Lvl 1	Inter-	Std	Lvl-1	V(τ)	
	Cnt	Cnt	Cnt	Cnt	Cnt	Cnt	Cnt	action	Dev	Units	/ σ^2	
Power	N	C	K00	K01	K10	K11	M	δ_{XZT}	σ	ρ_1	τ	Alpha
0.4191	500	5	5	5	5	5	5	3.00	9.80	0.100	0.100	0.050
0.6995	1000	5	10	10	10	10	5	3.00	9.80	0.100	0.100	0.050
0.8602	1500	5	15	15	15	15	5	3.00	9.80	0.100	0.100	0.050
0.9396	2000	5	20	20	20	20	5	3.00	9.80	0.100	0.100	0.050
0.6995	1000	10	5	5	5	5	5	3.00	9.80	0.100	0.100	0.050
0.9396	2000	10	10	10	10	10	5	3.00	9.80	0.100	0.100	0.050
0.9904	3000	10	15	15	15	15	5	3.00	9.80	0.100	0.100	0.050
0.9987	4000	10	20	20	20	20	5	3.00	9.80	0.100	0.100	0.050

Mixed Models Tests for Slope-Int'n in a 2x2 Fact. 3-Lvl Hier. Design with Rand. Slopes (Lvl-2 Rand.)

References

Ahn, C., Heo, M., and Zhang, S. 2015. Sample Size Calculations for Clustered and Longitudinal Outcomes in Clinical Research. CRC Press. New York.

Report Definitions


Power is the probability of rejecting a false null hypothesis. It should be close to one.
 N is the total number of level-1 units in the study.
 C is the number of level-2 units (subjects) per level-3 unit.
 K00 (Group 00 Count (X=0, Z=0)) is the number of level-2 units per level-3 unit in Group 00 (X=0 and Z=0).
 K01 (Group 01 Count (X=0, Z=1)) is the number of level-2 units per level-3 unit in Group 01 (X=0 and Z=1).
 K10 (Group 10 Count (X=1, Z=0)) is the number of level-2 units per level-3 unit in Group 10 (X=1 and Z=0).
 K11 (Group 11 Count (X=1, Z=1)) is the number of level-2 units per level-3 unit in Group 11 (X=1 and Z=1).
 M is the number of level-1 units per level-2 unit (i.e., the number of time points).
 δ_{xzt} is the 3-way interaction among the subject-specific slopes ($\beta_{11} - \beta_{10}$) - ($\beta_{01} - \beta_{00}$) at which the power is calculated.
 σ is the standard deviation of the Y_{ijk} assuming a fixed-slope model.
 ρ is the correlation among level-1 units in a particular level-2 unit.
 τ is ratio of the subject-specific slope variance, $V(\tau)$, to σ^2 .
 Alpha is the probability of rejecting a true null hypothesis, that is, rejecting when the means are actually equal.

Summary Statements

A total sample size of 500 observations, were obtained by sampling 5 level-3 units, and then sampling and then 5 level-2 units per level-3 unit in group 00, 5 level-2 units per level-3 unit in group 01, 5 level-2 units per level-3 unit in group 10, and 5 level-2 units per level-3 unit in group 11. Finally, 5 level-1 units (repeated measurements) were obtained from each level-2 unit. This sample achieves 42% power to detect a three-way interaction among the subject-specific slopes of at least 3.00. This 2x2 interaction was formed from the four group slopes across time using the following formula: $\delta_{xzt} = (\beta_{11} - \beta_{10}) - (\beta_{01} - \beta_{00})$. The standard deviation is 9.80. The intraclass correlation coefficient of level-1 units is 0.100. The ratio of the subject-specific slope variance to σ^2 is 0.100. A test based on a mixed-model analysis assuming random slopes will be used. This test will be conducted at a significance level of 0.050.

This report shows the power for each of the scenarios.

Plots Section



These plots show the power for the various parameter settings.

Example 2 – Calculating Sample Size (K00)

Continuing with the last example, suppose the researchers want to determine the value of K00 needed to achieve 90% power for both values of M.

Setup

This section presents the values of each of the parameters needed to run this example. First, from the PASS Home window, load the **Mixed Models Tests for Slope-Int'n in a 2x2 Fact. 3-Lvl Hier. Design with Rand. Slopes (Lvl-2 Rand.)** procedure window. You may then make the appropriate entries as listed below, or open **Example 2** by going to the **File** menu and choosing **Open Example Template**.

<u>Option</u>	<u>Value</u>
Design Tab	
Solve For	K00 (Group 00 Count (X=0, Z=0))
Power.....	0.90
Alpha.....	0.05
C (Level 3 Units)	5 10
K01 (Group 01 Count (X=0, Z=1))	K00
K10 (Group 10 Count (X=1, Z=0))	K00
K11 (Group 11 Count (X=1, Z=1))	K00
M (Level 1 Units Per Level-2 Unit).....	5
$\delta_{xz\tau}$ (Three-Way Interaction)	3
σ (Standard Deviation)	9.8
ρ_1 (Correlation Among Level-1 Units)	0.1
τ ($V(\tau) / \sigma^2$)	0.1

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results

Numeric Results for an Interaction in a 2x2 Factorial												
	Total Lvl 1 Cnt	Lvl 3 Cnt	Grp00 Lvl 2 Cnt	Grp01 Lvl 2 Cnt	Grp10 Lvl 2 Cnt	Grp11 Lvl 2 Cnt	Lvl 1 Cnt	2 x 2 Interaction $\delta_{xz\tau}$	Std Dev σ	ICC Lvl-1 Units ρ_1	$V(\tau) / \sigma^2$ τ	Alpha
Power 0.9149	1800	5	18	18	18	18	5	3.00	9.80	0.100	0.100	0.050
Power 0.9149	1800	10	9	9	9	9	5	3.00	9.80	0.100	0.100	0.050

This report shows the required K00 count for each of the scenarios.

Example 3 – Validation using Ahn, Heo, and Zhang (2015)

Ahn, Heo, and Zhang (2015) page 222 provide a table in which several scenarios are reported. We will validate this procedure by duplicating the results of the seventh row. Note that in using this table in this case, we reverse the second and third levels. Hence, instead of searching for C00, we will search for K00. Also, K in the table becomes C in our example.

The following parameter settings are used for the power analysis: Power = 0.80; $\delta_{XZT} = 0.3$; $\sigma = 4$; $\rho_1 = 0.1$; $\tau = 0.1$; $M = 5$; $C = 8$; and $\alpha = 0.05$. The values of K00, K01, K10, K11 are found to be 133 and the resulting power is 0.801.

Setup

This section presents the values of each of the parameters needed to run this example. First, from the PASS Home window, load the **Mixed Models Tests for Slope-Int'n in a 2x2 Fact. 3-Lvl Hier. Design with Rand. Slopes (Lvl-2 Rand.)** procedure window. You may then make the appropriate entries as listed below, or open **Example 3** by going to the **File** menu and choosing **Open Example Template**.

Option	Value
Design Tab	
Solve For	K00 (Group 00 Count (X=0, Z=0))
Power.....	0.80
Alpha.....	0.05
C (Level 3 Units)	8
K01 (Group 01 Count (X=0, Z=1))	K00
K10 (Group 10 Count (X=1, Z=0))	K00
K11 (Group 11 Count (X=1, Z=1))	K00
M (Level 1 Units Per Level-2 Unit).....	5
δ_{XZT} (Three-Way Interaction)	0.3
σ (Standard Deviation)	4
ρ_1 (Correlation Among Level-1 Units)	0.1
τ ($V(\tau) / \sigma^2$)	0.1

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results

Numeric Results for an Interaction in a 2x2 Factorial												
	Total Lvl 1 Cnt	Lvl 3 Cnt	Grp00 Lvl 2 Cnt	Grp01 Lvl 2 Cnt	Grp10 Lvl 2 Cnt	Grp11 Lvl 2 Cnt	Lvl 1 Cnt	2 x 2 Inter-action δ_{XZT}	Std Dev σ	ICC Lvl-1 Units ρ_1	$V(\tau) / \sigma^2$ τ	Alpha
Power	N	C	K00	K01	K10	K11	M					
0.8013	21280	8	133	133	133	133	5	0.30	4.00	0.100	0.100	0.050

PASS also calculates K00 to be 133 and the power to be 0.8013.