

Chapter 331

Multi-Arm Tests for the Ratio of Treatment and Control Means (Normal Data)

Introduction

This module computes power and sample size for multiple comparisons of treatment means versus a control mean when the data are assumed to follow the normal distribution and the statistical hypotheses are expressed in terms of mean ratios. Note that when the data follow a log-normal distribution rather than the normal distribution so that a log transformation is used, you should use other **PASS** procedures that assume a log-normal data distribution.

The details of this t-test are given in Rothmann, Wiens, and Chan (2012) and, to lesser extent, in Kieser and Hauschke (1999). The multiple comparison aspect of this procedure is based on the results in Machin, Campbell, Tan, and Tan (2018).

In this parallel-group design, there are k treatment groups and one control group. A mean is measured in each group. A total of k hypothesis tests are anticipated each comparing a treatment group with the common control group using a t-test based on the ratio of two means.

The Bonferroni adjustment of the type I error rate may be optionally made because several comparisons are being tested using the same data. Making a multiplicity adjustment is usually recommended, but not always. In fact, Saville (1990) advocates not applying it and Machin, Campbell, Tan, and Tan (2018) include omitting it as a possibility.

Background

Whether you want to test several doses of a single treatment or several types of treatments, good research practice requires that each treatment be compared with a control. A popular three-arm design consists of three groups: control, treatment A, and treatment B. Two tests are run: treatment A versus control and treatment B versus the same control. This design avoids having to obtain a second control group for treatment B. Besides the obvious efficiency in subjects, it may be easier to recruit subjects if their chances of receiving the new treatment are better than 50-50.

Technical Details

Statistical Hypotheses Based on Mean Ratios

Suppose you want to compare k treatment groups with means μ_i and sample sizes N_i and one control group with mean μ_C and sample size N_C . The total sample size is $N = N_1 + N_2 + \dots + N_k + N_C$.

Let $R = \mu_i/\mu_C$ and $R_0 = \mu_{i|H_0}/\mu_C$. For each treatment, the null and alternative two-sided hypotheses are

$$H_0: R = R_0 \quad \text{vs.} \quad H_1: R \neq R_0.$$

The one-sided hypotheses are

$$H_0: R \leq R_0 \quad \text{vs.} \quad H_1: R > R_0$$

and

$$H_0: R \geq R_0 \quad \text{vs.} \quad H_1: R < R_0.$$

In these formulas, usually R_0 is set to 1, but this is not always the case.

Test Statistic – Equal Variance T-Test

A suitable Type I error probability is chosen for the tests, the data is collected, and a t -statistic is generated for each treatment using the formula

$$t = \frac{\bar{x}_i - R_0 \bar{x}_C}{\sqrt{\frac{(N_i - 1)s_i^2 + (N_C - 1)s_C^2}{N_i + N_C - 2} \left(\frac{1}{N_i} + \frac{R_0^2}{N_C} \right)}}$$

where \bar{X}_i and \bar{X}_C are the sample means of the treatment and control groups and the denominator uses the pooled estimate of the standard deviation. This t -statistic follows a central t distribution with $N_i + N_C - 2$ degrees of freedom.

Power Calculation

For a specified alternative $R = R_1$, t follows the noncentral t distribution with $N_1 + N_2 - 2$ degrees of freedom and noncentrality

$$\left(\frac{R - R_0}{\sigma/\mu_c} \right) \sqrt{\frac{N_C}{\frac{N_C}{N_i} + R_0^2}}$$

Hence, the power of a one-sided test is given by

$$(1 - \beta) = \Pr(t \geq t_{1-\alpha, N_i + N_C - 2} | R, R_0, \sigma)$$

The power of the two-sided test is calculated similarly.

Multiplicity Adjustment

Because k t -tests between treatment groups and the control group are run when analyzing the results of this study, many statisticians recommend that the Bonferroni adjustment be applied. This adjustment is easy to apply: the value of alpha that is used in the test is found by dividing the original alpha by the number of tests. For example, if the original alpha is set at 0.05 and the number of treatment (not including the control) groups is five, the individual tests will be conducted using an alpha of 0.01.

The main criticism of this procedure is that if there are many tests, the value of alpha becomes very small. To mitigate against this complaint, some statisticians recommend separating the treatment groups into those that are of primary interest and those that are of secondary interest. The Bonferroni adjustment is made by using the number of primary treatments rather than the total number of treatments.

There are some who advocate ignoring the adjustment entirely in the case of randomized clinical trials. See for example Saville (1990) and the discussion in chapter 14 of Machin, Campbell, Tan, and Tan (2018).

Size of the Control Group

Because the control group is used over and over, some advocate increasing the number of subjects in this group. The standard adjustment is to include \sqrt{k} subjects in the control group for each subject in one of the treatment groups. See Machin, Campbell, Tan, and Tan (2018, pages 231-232). Note that often, the treatment groups all have the same size.

Example 1 – Finding the Sample Size

A parallel-group clinical trial is being designed to compare three treatment therapies against the standard therapy. Lower values of the response is desirable. Suppose the standard therapy has mean response of 9.3 with a standard deviation of 2.5.

The investigators would like a sample size large enough to find statistical significance at the 0.05 level if the actual mean responses of the three treatments are 7.3, 7.6, and 8.1 and the power is 0.80 in each test. They want to consider a range of standard deviations from 2.0 to 3.0. The tests will be two-sided.

Following standard procedure, the control group multiplier will be set to $\sqrt{k} = \sqrt{3} = 1.732$ since the control group is used for three comparisons in this design.

Setup

If the procedure window is not already open, use the PASS Home window to open it. The parameters for this example are listed below and are stored in the **Example 1** settings file. To load these settings to the procedure window, click **Open Example Settings File** in the Help Center or File menu.

Design Tab

| | |
|-------------------------------------|---|
| Solve For | Sample Size |
| Alternative Hypothesis | Two-Sided (H1: R ≠ R0) |
| Power of Each Test | 0.8 |
| Overall Alpha | 0.05 |
| Bonferroni Adjustment | Standard Bonferroni |
| Group Allocation | Enter Group Allocation Pattern, solve for group sample sizes |
| R0 (Mean Ratio H0) | 1 |
| Control Mean | 9.3 |
| Control Sample Size Allocation..... | 1.732 |
| Set A Number of Groups..... | 1 |
| Set A Mean | 7.3 |
| Set A Sample Size Allocation | 1 |
| Set B Number of Groups..... | 1 |
| Set B Mean | 7.6 |
| Set B Sample Size Allocation | 1 |
| Set C Number of Groups | 1 |
| Set C Mean..... | 8.1 |
| Set C Sample Size Allocation | 1 |
| Set D Number of Groups | 0 |
| More..... | Unchecked |
| σ (Standard Deviation)..... | 2 2.5 3 |

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Reports

Numeric Results

Solve For: **Sample Size**
 Group Allocation: Enter Group Allocation Pattern, solve for group sample sizes
 Test Type: Equal Variances T-Test
 Hypotheses: H0: R = R0 vs. H1: R ≠ R0
 Number of Groups: 4
 Bonferroni Adjustment: Standard Bonferroni (Divisor = 3)

| Comparison | Power | | Sample Size | | Mean μ_i | Mean Ratio | | Standard Deviation σ | COVi | Alpha | |
|------------|--------|---------|-------------|------------|--------------|-------------|-------------|-----------------------------|---------|---------|---------------------|
| | Target | Actual | Ni | Allocation | | Ratio H0 R0 | Ratio H1 Ri | | | Overall | Bonferroni-Adjusted |
| Control | | | 83 | 1.732 | 9.3 | | | 2.0 | 0.21505 | | |
| vs A | 0.8 | 0.99889 | 48 | 1.000 | 7.3 | 1 | 0.78495 | 2.0 | 0.27397 | 0.05 | 0.01667 |
| vs B | 0.8 | 0.98749 | 48 | 1.000 | 7.6 | 1 | 0.81720 | 2.0 | 0.26316 | 0.05 | 0.01667 |
| vs C | 0.8 | 0.81003 | 48 | 1.000 | 8.1 | 1 | 0.87097 | 2.0 | 0.24691 | 0.05 | 0.01667 |
| Total | | | 227 | | | | | | | | |
| Control | | | 126 | 1.732 | 9.3 | | | 2.5 | 0.26882 | | |
| vs A | 0.8 | 0.99867 | 73 | 1.000 | 7.3 | 1 | 0.78495 | 2.5 | 0.34247 | 0.05 | 0.01667 |
| vs B | 0.8 | 0.98593 | 73 | 1.000 | 7.6 | 1 | 0.81720 | 2.5 | 0.32895 | 0.05 | 0.01667 |
| vs C | 0.8 | 0.80111 | 73 | 1.000 | 8.1 | 1 | 0.87097 | 2.5 | 0.30864 | 0.05 | 0.01667 |
| Total | | | 345 | | | | | | | | |
| Control | | | 182 | 1.732 | 9.3 | | | 3.0 | 0.32258 | | |
| vs A | 0.8 | 0.99873 | 105 | 1.000 | 7.3 | 1 | 0.78495 | 3.0 | 0.41096 | 0.05 | 0.01667 |
| vs B | 0.8 | 0.98633 | 105 | 1.000 | 7.6 | 1 | 0.81720 | 3.0 | 0.39474 | 0.05 | 0.01667 |
| vs C | 0.8 | 0.80333 | 105 | 1.000 | 8.1 | 1 | 0.87097 | 3.0 | 0.37037 | 0.05 | 0.01667 |
| Total | | | 497 | | | | | | | | |

- Comparison: The group that is involved in the comparison between the treatment and control displayed on this report line. The comparison is made using the ratio.
- Target Power: The power desired. Power is probability of rejecting a false null hypothesis for this comparison. This power is of the comparison shown on this line only.
- Actual Power: The power actually achieved.
- Ni: Sample Size. The number of subjects in the ith group. The total sample size, N, is shown as the last row of the column.
- Allocation: The group sample size allocation pattern. The value on each row represents the relative number of subjects assigned to the group.
- μ_i : The mean of the ith group at which the power is computed. The first row contains μ_c , the control group mean.
- R0: The ratio of means assumed by the null hypothesis. $R0 = (\mu_i|H0) / \mu_c$.
- Ri: The ratio of the means at which the power is calculated. $Ri = \mu_i / \mu_c$.
- σ : The standard deviation of the responses within each group.
- COVi: The coefficient of variation of the ith group. $COVi = \sigma / \mu_i$.
- Overall Alpha: The probability of rejecting at least one of the comparisons in this experiment when each null hypothesis is true.
- Bonferroni Alpha: The adjusted significance level at which each individual comparison is made.

Summary Statements

A parallel, 4-group design (with one control group and 3 treatment groups) will be used to test whether the mean for each treatment group is different from the control group mean, with a null mean ratio of 1 (H0: R = 1 versus H1: R ≠ 1, R = μ_i / μ_c). The hypotheses will be evaluated using 3 two-sided, two-sample, Bonferroni-adjusted, equal-variance, ratio-based t-tests, with an overall (experiment-wise) Type I error rate (α) of 0.05. The common standard deviation for all groups is assumed to be 2. The control group mean is assumed to be 9.3. To detect the treatment means 7.3, 7.6, and 8.1 with at least 80% power for each test, the control group sample size needed will be 83 and the number of needed subjects for the treatment groups will be 48, 48, and 48 (totaling 227 subjects overall).

Multi-Arm Tests for the Ratio of Treatment and Control Means (Normal Data)

Dropout-Inflated Sample Size

| Group | Dropout Rate | Sample Size Ni | Dropout- Inflated Enrollment Sample Size Ni' | Expected Number of Dropouts Di |
|-------|--------------|-------------------|--|---|
| 1 | 20% | 83 | 104 | 21 |
| 2 | 20% | 48 | 60 | 12 |
| 3 | 20% | 48 | 60 | 12 |
| 4 | 20% | 48 | 60 | 12 |
| Total | | 227 | 284 | 57 |
| 1 | 20% | 126 | 158 | 32 |
| 2 | 20% | 73 | 92 | 19 |
| 3 | 20% | 73 | 92 | 19 |
| 4 | 20% | 73 | 92 | 19 |
| Total | | 345 | 434 | 89 |
| 1 | 20% | 182 | 228 | 46 |
| 2 | 20% | 105 | 132 | 27 |
| 3 | 20% | 105 | 132 | 27 |
| 4 | 20% | 105 | 132 | 27 |
| Total | | 497 | 624 | 127 |

| | |
|--------------|---|
| Group | Lists the group numbers. |
| Dropout Rate | The percentage of subjects (or items) that are expected to be lost at random during the course of the study and for whom no response data will be collected (i.e., will be treated as "missing"). Abbreviated as DR. |
| Ni | The evaluable sample size for each group at which power is computed (as entered by the user). If Ni subjects are evaluated out of the Ni' subjects that are enrolled in the study, the design will achieve the stated power. |
| Ni' | The number of subjects that should be enrolled in each group in order to obtain Ni evaluable subjects, based on the assumed dropout rate. Ni' is calculated by inflating Ni using the formula $Ni' = Ni / (1 - DR)$, with Ni' always rounded up. (See Julious, S.A. (2010) pages 52-53, or Chow, S.C., Shao, J., Wang, H., and Lakhnygina, Y. (2018) pages 32-33.) |
| Di | The expected number of dropouts in each group. $Di = Ni' - Ni$. |

Dropout Summary Statements


Anticipating a 20% dropout rate, group sizes of 104, 60, 60, and 60 subjects should be enrolled to obtain final group sample sizes of 83, 48, 48, and 48 subjects.

References

- Rothmann, M.D., Wiens, B.L., and Chan, I.S.F. 2012. Design and Analysis of Non-Inferiority Trials. Taylor & Francis/CRC Press. Boca Raton, Florida.
- Kieser, M. and Hauschke, D. 1999. 'Approximate Sample Sizes for Testing Hypotheses about the Ratio and Difference of Two Means.' Journal of Biopharmaceutical Studies, Volume 9, No. 4, pages 641-650.
- Julious, Steven A. 2004. 'Tutorial in Biostatistics. Sample sizes for clinical trials with Normal data.' Statistics in Medicine, 23:1921-1986.
- Chow, S.C., Shao, J., Wang, H., and Lakhnygina, Y. 2018. Sample Size Calculations in Clinical Research, 3rd Edition. Chapman & Hall/CRC. Boca Raton, FL. Pages 86-88.
- Machin, D., Campbell, M.J., Tan, S.B, and Tan, S.H. 2018. Sample Sizes for Clinical, Laboratory, and Epidemiology Studies, 4th Edition. Wiley Blackwell.

This report shows the numeric results of this power study. Notice that the results are shown in blocks of three rows at a time. Each block represents a single design.

Plots Section



This plot gives a visual presentation to the results in the Numeric Report. We can quickly see the impact on the sample size of changing the standard deviation.

Example 2 – Validation using a Previously Validated Procedure

We could not find a validation result in the statistical literature, so we will use a previously validated **PASS** procedure (**Tests for the Ratio of Two Means (Normal Data)**) to produce the results for the following example.

A parallel-group clinical trial is being designed to compare three treatment therapies against the standard therapy. Suppose the standard therapy has mean response of 9.3 with a standard deviation of 2.5. This standard deviation results in a COV of 0.26882. The investigators would like a sample size large enough to find statistical significance at the $0.05/3 = 0.01667$ level if the actual mean responses of the three treatments are 7.3, 7.6, and 8.1 and the power is 0.80 in each test. These three means result in R1 values of 0.78495, 0.81720, and 0.87097. The tests will be two-sided.

The sample sizes of all groups will be equal.

The **Tests for the Ratio of Two Means (Normal Data)** procedure is set up as follows.

| Design Tab | |
|--|-------------------------------------|
| Solve For | Sample Size |
| Alternative Hypothesis | Two-Sided (H1: R ≠ R0) |
| Test Type..... | Equal Variances T-Test |
| Power..... | 0.8 |
| Alpha..... | 0.01667 (which is Alpha / k) |
| Group Allocation | Equal (N1 = N2) |
| R0 (Ratio H0)..... | 1 |
| R1 (Ratio H1)..... | 0.78495 0.81720 0.87097 |
| CV (Coef of Variation, σ^2 / μ^2)..... | 0.26886 |

This set of options generates the following report.

| Numeric Results | | | | | | | | |
|-----------------|---------------------------|----------------|----|-----|------------------------------|---------|--|--|
| Solve For: | Sample Size | | | | | | | |
| Ratio: | $R = \mu_1 / \mu_2$ | | | | | | | |
| Hypotheses: | H0: R = R0 vs. H1: R ≠ R0 | | | | | | | |
| Test: | Equal Variances T-Test | | | | | | | |
| Target Power | Actual Power | Ratio of Means | | | Coef of Variation Control CV | Alpha | | |
| | | N1 | N2 | N | | | | |
| 0.8 | 0.81092 | 35 | 35 | 70 | 0.26886 | 0.01667 | | |
| 0.8 | 0.80249 | 47 | 47 | 94 | 0.26886 | 0.01667 | | |
| 0.8 | 0.80321 | 93 | 93 | 186 | 0.26886 | 0.01667 | | |

In order to maintain a power of 80% for all three groups, it is apparent that the groups will all need to have a sample size of 93. This table contains the validation values. We will now run these values through the current procedure and compare the results with these values.

Setup

If the procedure window is not already open, use the PASS Home window to open it. The parameters for this example are listed below and are stored in the **Example 2** settings file. To load these settings to the procedure window, click **Open Example Settings File** in the Help Center or File menu.

| Design Tab | |
|------------------------------|-----------------------------------|
| Solve For | Sample Size |
| Alternative Hypothesis | Two-Sided (H1: R ≠ R0) |
| Power of Each Test | 0.8 |
| Overall Alpha | 0.05 |
| Bonferroni Adjustment | Standard Bonferroni |
| Group Allocation | Equal (Nc = N1 = N2 = ...) |
| R0 (Mean Ratio H0) | 1 |
| Control Mean | 9.3 |
| Set A Number of Groups..... | 1 |
| Set A Mean | 7.3 |
| Set B Number of Groups..... | 1 |
| Set B Mean | 7.6 |
| Set C Number of Groups | 1 |
| Set C Mean..... | 8.1 |
| Set D Number of Groups | 0 |
| More..... | Unchecked |
| σ (Standard Deviation)..... | 2.5 |

Output

Click the Calculate button to perform the calculations and generate the following output.

| Numeric Results | | | | | | | | | | |
|------------------------|--------|-----------------------------------|----------------|---------|-------------|-------------|----------------------|---------|---------|---------------------|
| Solve For: | | Sample Size | | | | | | | | |
| Group Allocation: | | Equal (Nc = N1 = N2 = ...) | | | | | | | | |
| Test Type: | | Equal Variances T-Test | | | | | | | | |
| Hypotheses: | | H0: R = R0 vs. H1: R ≠ R0 | | | | | | | | |
| Number of Groups: | | 4 | | | | | | | | |
| Bonferroni Adjustment: | | Standard Bonferroni (Divisor = 3) | | | | | | | | |
| Comparison | Power | | Sample Size Ni | Mean μi | Mean Ratio | | Standard Deviation σ | COVi | Alpha | |
| | Target | Actual | | | Ratio H0 R0 | Ratio H1 Ri | | | Overall | Bonferroni-Adjusted |
| Control | | | 93 | 9.3 | | | 2.5 | 0.26882 | | |
| vs A | 0.8 | 0.99873 | 93 | 7.3 | 1 | 0.78495 | 2.5 | 0.34247 | 0.05 | 0.01667 |
| vs B | 0.8 | 0.98633 | 93 | 7.6 | 1 | 0.81720 | 2.5 | 0.32895 | 0.05 | 0.01667 |
| vs C | 0.8 | 0.80335 | 93 | 8.1 | 1 | 0.87097 | 2.5 | 0.30864 | 0.05 | 0.01667 |
| Total | | | 372 | | | | | | | |

As you can see, the sample sizes are all 93, which match the largest sample size found in the validation run above. The procedure is validated.