

Chapter 161

Non-Inferiority Tests for the Ratio of Two Correlated Proportions

Introduction

This module computes power and sample size for non-inferiority tests of the ratio in which two dichotomous responses are measured on each subject. When one is interested in showing that the true proportions are different, the data are often analyzed with McNemar's test. However, we are interested in showing non-inferiority rather than difference. For example, suppose a diagnostic procedure is accurate, but is expensive to apply or has serious side effects. A replacement procedure is sought which is no less accurate, but is less expensive or has fewer side effects. In this case, we are not interested in showing that the two diagnostic procedures are different, but rather that the second is no worse than the first. *Non-inferiority tests* were designed for this situation.

These tests are often divided into two categories: *equivalence* (two-sided) tests and *non-inferiority* (one-sided) tests. Here, the term *equivalence tests* means that we want to show that two diagnostic procedures are equivalent—that is, their accuracy is about the same. This requires a two-sided hypothesis test. On the other hand, *non-inferiority tests* are used when we want to show that a new (experimental) procedure is no worse than the existing (reference or gold-standard) one. This requires a one-sided hypothesis test. The procedures discussed in this chapter deal with the non-inferiority (one-sided) case.

Technical Details

The results of a study in which two dichotomous responses are measured on each subject can be displayed in a 2-by-2 table in which one response is shown across the columns and the other is shown down the rows. In the discussion to follow, the columns of the table represent the standard (reference or control) response and the rows represent the treatment (experimental) response. The outcome probabilities can be classified into the following table.

Experimental Diagnosis	Standard Yes	Diagnosis No	Total
Yes	p_{11}	p_{10}	P_T
No	p_{01}	p_{00}	$1 - P_T$
Total	P_S	$1 - P_S$	1

Non-Inferiority Tests for the Ratio of Two Correlated Proportions

In this table, $p_{ij} = p_{Treatment, Standard}$. That is, the first subscript represents the response of the new, experimental procedure while the second subscript represents the response of the standard procedure. Thus, p_{01} represents the proportion having a negative treatment response and a positive standard response.

Sensitivity, Specificity, and Prevalence

To aid in interpretation, analysts have developed a few proportions that summarize the table. Three of the most popular ratios are *sensitivity*, *specificity*, and *prevalence*.

Sensitivity

Sensitivity is the proportion of subjects with a positive standard response who also have a positive experimental response. In terms of proportions from a 2-by-2 table,

$$\text{Sensitivity} = \frac{p_{11}}{(p_{01} + p_{11})} = \frac{p_{11}}{P_S}$$

Specificity

Specificity is the proportion of subjects with a negative standard response who also have a negative experimental response. In terms of proportions from a 2-by-2 table,

$$\text{Specificity} = \frac{p_{00}}{(p_{10} + p_{00})}$$

Prevalence

Prevalence is the overall proportion of individuals with the disease (or feature of interest). In terms of proportions from a 2-by-2 table,

$$\text{Prevalence} = P_S$$

Table Probabilities

The outcome counts from a sample of n subjects can be classified into the following table.

Experimental Diagnosis	Standard Diagnosis		Total
	Yes	No	
Yes	n_{11}	n_{10}	n_T
No	n_{01}	n_{00}	$n - n_T$
Total	n_S	$n - n_S$	n

Note that $n_{11} + n_{00}$ is the number of matches (*concordant pairs*) and $n_{01} + n_{10}$ is the number of *discordant pairs*.

The hypothesis of interest concerns the two marginal probabilities P_T and P_S . P_S represents the accuracy or success of the standard test and P_T represents the accuracy or success of the new, experimental test. Non-inferiority is defined in terms of either the difference of these two proportions, $D = P_T - P_S$, or the relative risk ratio, $R = P_T/P_S$. The choice between D and R will usually lead to different sample sizes to achieve the same power.

Non-Inferiority Hypotheses using Ratios

The following is based on Nam and Blackwelder (2002). We refer you to this paper for the complete details of which we will only provide a brief summary here.

If we define R_0 as the non-inferiority ratio with $0 < R_0 < 1$, then the null and alternative hypotheses of non-inferiority in terms of the ratio are

$$H_0: P_T/P_S \leq R_0 \quad \text{versus} \quad H_1: P_T/P_S > R_0.$$

Test Statistics

The test statistic for an asymptotic test based on constrained maximum likelihood for large n is given by

$$Z(R_0) = \frac{\sqrt{n(\hat{P}_T - R_0\hat{P}_S)}}{\sqrt{R_0(\tilde{p}_{10} + \tilde{p}_{01})}}$$

where

$$\tilde{p}_{10} = \frac{-\hat{P}_T + R_0^2(\hat{P}_S + 2\hat{p}_{10}) + \sqrt{(\hat{P}_T - R_0^2\hat{P}_S)^2 + 4R_0^2\hat{p}_{10}\hat{p}_{01}}}{2R_0(R_0 + 1)}$$

$$\tilde{p}_{01} = R_0\tilde{p}_{10} - (R_0 - 1)(1 - \hat{p}_{00})$$

$$\hat{p}_{01} = \frac{n_{01}}{n}, \hat{p}_{10} = \frac{n_{10}}{n}, \hat{P}_T = \frac{n_{10} + n_{11}}{n}, \hat{P}_S = \frac{n_{01} + n_{11}}{n}$$

Power Formula

The power when the true value of the relative risk ratio is R_1 can be evaluated exactly using the multinomial distribution. When n is large, we use a normal approximation to the multinomial distribution which leads to

$$\text{Power} = 1 - \Phi(c_{NI})$$

where

$$c_{NI} = \frac{z_{1-\alpha}\sqrt{\bar{V}_0(R_0)} - E_1(R_0)}{\sqrt{V_1(R_0)}}$$

$$\bar{V}_0(R_0) = \frac{R_0(\bar{p}_{10} + \bar{p}_{01})}{n}$$

$$E_1(R_0) = P_S(R_1 - R_0)$$

$$V_1(R_0) = \frac{P_S(R_1 + R_0^2) - 2R_0p_{11} - P_S^2(R_1 - R_0)^2}{n}$$

$$\bar{p}_{10} = \frac{-P_T + R_0^2(P_S + 2p_{10}) + \sqrt{(P_T - R_0^2P_S)^2 + 4R_0^2p_{10}p_{01}}}{2R_0(R_0 + 1)}$$

$$\bar{p}_{01} = R_0\bar{p}_{10} - (R_0 - 1)(1 - p_{00})$$

Nuisance Parameter

The 2-by-2 table includes four parameters, p_{11} , p_{10} , p_{01} , and p_{00} , but the power calculations only require two parameters: P_S and R_1 . A third parameter is defined implicitly since the sum of the four parameters is one. Thus, one parameter (known as a nuisance parameter) remains unaccounted for. This parameter must be addressed to fully specify the problem. This fourth parameter can be specified using any one of the following: p_{11} , p_{10} , p_{01} , p_{00} , $p_{10} + p_{01}$, $p_{11} + p_{00}$, or the sensitivity of the experimental response, p_{11}/P_S .

It may be difficult to specify a reasonable value for the nuisance parameter since its value may not be even approximately known until after the study is conducted. Because of this, we suggest that you calculate power or sample size for a range of values of the nuisance parameter. This will allow you to determine how sensitive the results are to its value.

Procedure Options

This section describes the options that are specific to this procedure. These are located on the Design tab. For more information about the options of other tabs, go to the Procedure Window chapter.

Design Tab

The Design tab contains the parameters associated with this test such as the proportions, sample sizes, alpha, and power.

Solve For

Solve For

This option specifies the parameter to be solved for from the other parameters. The parameters that may be selected are *Power* or *Sample Size*.

Power Calculation

Power Calculation Method

Select the method to be used to calculate power.

The choices are

- **Multinomial Enumeration**

Power is computed using multinomial enumeration of all possible outcomes when $N \leq \text{Max } N$ for Multinomial Enumeration (otherwise, the normal approximation is used). Multinomial enumeration of all outcomes is possible because of the discrete nature of the data.

- **Normal Approximation**

Approximate power is computed using the normal approximation to the multinomial distribution.

The exact calculation using the multinomial distribution becomes very time consuming for $N > 500$. When $N > 500$, the difference between the multinomial and approximate calculations is small.

For small values of N (less than 100), the Multinomial Enumeration power may be overly optimistic because the discrete nature of the trinomial distribution results in the actual alpha value being higher than its target. To be on the safe side, we recommend that you use the approximate calculation.

Non-Inferiority Tests for the Ratio of Two Correlated Proportions

Max N for Multinomial Enumeration

Only shown when Power Calculation Method = "Multinomial Enumeration"

Specify the maximum value of N (sample size) that uses the exact power calculation based on the multinomial distribution. N's greater than this value will use the asymptotic approximation.

Power and Alpha

Power

This option specifies one or more values for power. Power is the probability of rejecting a false null hypothesis, and is equal to one minus Beta. Beta is the probability of a type-II error, which occurs when a false null hypothesis is not rejected. Here, a type-II error occurs when you fail to conclude non-inferiority when in fact it is true.

Values must be between zero and one. Historically, the value of 0.80 (Beta = 0.20) was used for power. Now, 0.90 (Beta = 0.10) is also commonly used.

A single value may be entered here or a range of values such as *0.8 to 0.95 by 0.05* may be entered.

Alpha

This option specifies one or more values for the probability of a type-I error. A type-I error occurs when a true null hypothesis is rejected. Here, a type-I error occurs when you falsely conclude non-inferiority.

Sample Size

N (Sample Size)

Enter a value for the sample size. This value must be greater than two. You may enter a range of values such as *10 to 100 by 10*.

Effect Size – Ratios

R0 (Non-Inferiority Ratio)

Specify the ratio, P_t/P_s , for which the treatment group will be considered non-inferior to the standard group. You can enter a single value such as *0.8* or a series of values such as *0.8 0.85 0.9* or *0.8 to 0.9 by 0.05* in the range $0 < R_0 < 1$ and $R_0 < R_1$.

R1 (Actual Ratio)

Specify the actual ratio, P_t/P_s , at which power will be calculated. This value is used to calculate P_t using the formula

$$P_t = P_s \times D_1.$$

Often this value is set equal to 1, but this is not required. Enter values in the range $R_1 > R_0$.

Be careful that the value you enter here along with other values in this procedure do not result in an undefined 2×2 table. The values entered in this procedure must result in a 2×2 table whose cell values and marginal totals each sum to 1.

Non-Inferiority Tests for the Ratio of Two Correlated Proportions

Effect Size – Standard Proportion

Ps (Standard Proportion)

The proportion of Yes's (positive responses or successes) when subjects receive the standard procedure.

You may enter a set of values separated by blanks. For example, you could enter *0.50 0.60 0.70*. Enter values in the range $0 < P_s < 1$.

Be careful that the value you enter here along with other values in this procedure do not result in an undefined 2×2 table. The values entered in this procedure must result in a 2×2 table whose cell values and marginal totals each sum to 1.

Effect Size – Nuisance Parameter

Nuisance Parameter Type

Enter the type of nuisance parameter here. Unfortunately, the 2-by-2 table cannot be completely specified by using only the parameters P_s and D_a or P_s and R_a . One other parameter must be specified. This additional parameter is called a “nuisance” parameter. It will be assumed to be a known quantity. Several possible choices are available. This option lets you specify which parameter you want to use. In all cases, the value you specify is a proportion.

- **P11 (% Positive Matches)**
The proportion of subjects that are positive on both tests.
- **P00 (% Negative Matches)**
The proportion of subjects that are negative on both tests.
- **P01 (% -Trt +Std)**
The proportion of subjects that are negative on the treatment, but positive on the standard.
- **P10 (% +Trt -Std)**
The proportion of subjects that are positive on the treatment, but negative on the standard.
- **P11+P00 (% Matches)**
The proportion of matches (concordant pairs).
- **P01+P10 (% Disagree)**
The proportion of non-matches (discordant pairs).
- **P11/Ps (Sensitivity)**
The sensitivity.

Nuisance Parameter Value

Enter the value of the nuisance parameter that you specified in the “Nuisance Parameter Type” box. This value is a proportion, so it must be between 0 and 1.

Non-Inferiority Tests for the Ratio of Two Correlated Proportions

Example 1 – Finding Power

Researchers have developed a new treatment for migraine headaches which is less expensive than a current standard. The researchers need to show that the proportion of individuals who respond to the new treatment is not inferior to the standard treatment. The new treatment will be considered non-inferior if its success rate is no less than 95% of the success rate of the standard, which is about 0.65. They want to study the power for various sample sizes between 500 and 4000 at the 5% significance level. They'll study various values of the nuisance parameter: $P_{11}/P_s = \text{sensitivity}$ (0.5 to 0.9).

Setup

This section presents the values of each of the parameters needed to run this example. First, from the PASS Home window, load the **Non-Inferiority Tests for the Ratio of Two Correlated Proportions** procedure window by expanding **Proportions**, then **Two Correlated Proportions**, then clicking on **Non-Inferiority**, and then clicking on **Non-Inferiority Tests for the Ratio of Two Correlated Proportions**. You may then make the appropriate entries as listed below, or open **Example 1** by going to the **File** menu and choosing **Open Example Template**.

<u>Option</u>	<u>Value</u>
Design Tab	
Solve For	Power
Power Calculation Method.....	Normal Approximation
Alpha.....	0.05
N (Sample Size).....	500 to 4000 by 500
R0 (Non-Inferiority Ratio).....	0.95
R1 (Actual Ratio)	1.0
Ps (Standard Proportion)	0.65
Nuisance Parameter Type	P11/Ps (Sensitivity)
Nuisance Parameter Value.....	0.5 to 0.9 by 0.1

Annotated Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results

Numeric Results

Hypotheses: $H_0: P_t/P_s \leq R_0$ vs. $H_1: P_t/P_s > R_0$

Power	Sample Size N	Non-Inf. Ratio R0	Actual Ratio R1	Treatment Proportion Pt	Standard Proportion Ps	Nuisance Parameter P11/Ps	Alpha
0.23552	500	0.950	1.000	0.650	0.650	0.500	0.050
0.27009	500	0.950	1.000	0.650	0.650	0.600	0.050
0.32464	500	0.950	1.000	0.650	0.650	0.700	0.050
0.42349	500	0.950	1.000	0.650	0.650	0.800	0.050
0.64700	500	0.950	1.000	0.650	0.650	0.900	0.050
0.36772	1000	0.950	1.000	0.650	0.650	0.500	0.050
0.42680	1000	0.950	1.000	0.650	0.650	0.600	0.050
0.51572	1000	0.950	1.000	0.650	0.650	0.700	0.050
0.65984	1000	0.950	1.000	0.650	0.650	0.800	0.050
0.89100	1000	0.950	1.000	0.650	0.650	0.900	0.050
0.48241	1500	0.950	1.000	0.650	0.650	0.500	0.050
0.55722	1500	0.950	1.000	0.650	0.650	0.600	0.050
0.66224	1500	0.950	1.000	0.650	0.650	0.700	0.050

(report continues)

* Power was computed using the normal approximation method.

Non-Inferiority Tests for the Ratio of Two Correlated Proportions

Report Definitions


Power is the probability of rejecting the null hypothesis when it is false.
 N is the number of subjects, the sample size.
 R0 is the ratio at which the treatment will be considered non-inferior to the standard.
 $R1 = P_t/P_s$ is the actual ratio at which the power is calculated.
 P_t is the response proportion in the treatment (experimental) group.
 P_s is the response proportion in the standard (baseline, reference, or control) group.
 The Nuisance Parameter is a value that is needed, but is not a direct part of the hypotheses.
 Alpha is the probability of rejecting the null hypothesis when it is true.

Summary Statements

For a one-sided non-inferiority test of the ratio of two correlated proportions, a sample size of 500 subjects achieves 23.552% power at a significance level of 0.050 when the non-inferiority ratio is 0.950, the treatment proportion is 0.650, the standard proportion is 0.650, and the actual ratio of the proportions is 1.000. These results assume that P_{11}/P_s is equal to 0.500.

This report shows the power for the indicated scenarios. All of the columns are defined in the 'Report Definitions' section.

Plots Section



These plots show the power versus the sample size for the various values of sensitivity. In this example, we see that the value of the nuisance parameter has a large effect on the calculated power.

Non-Inferiority Tests for the Ratio of Two Correlated Proportions

Example 2 – Finding Sample Size

Continuing with Example 1, the analysts want to determine the exact sample size necessary to achieve 90% power for all values of the nuisance parameter.

Setup

This section presents the values of each of the parameters needed to run this example. First, from the PASS Home window, load the **Non-Inferiority Tests for the Ratio of Two Correlated Proportions** procedure window by expanding **Proportions**, then **Two Correlated Proportions**, then clicking on **Non-Inferiority**, and then clicking on **Non-Inferiority Tests for the Ratio of Two Correlated Proportions**. You may then make the appropriate entries as listed below, or open **Example 2** by going to the **File** menu and choosing **Open Example Template**.

<u>Option</u>	<u>Value</u>
Design Tab	
Solve For	Sample Size
Power Calculation Method.....	Normal Approximation
Power.....	0.90
Alpha.....	0.05
R0 (Non-Inferiority Ratio).....	0.95
R1 (Actual Ratio)	1.0
Ps (Standard Proportion)	0.65
Nuisance Parameter Type	P11/Ps (Sensitivity)
Nuisance Parameter Value.....	0.5 to 0.9 by 0.1

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results							
Hypotheses: $H_0: P_t/P_s \leq R_0$ vs. $H_1: P_t/P_s > R_0$							
	Sample Size N	Non-Inf. Ratio R0	Actual Ratio R1	Treatment Proportion Pt	Standard Proportion Ps	Nuisance Parameter P11/Ps	Alpha
0.90004	5013	0.950	1.000	0.650	0.650	0.500	0.050
0.90001	4013	0.950	1.000	0.650	0.650	0.600	0.050
0.90004	3015	0.950	1.000	0.650	0.650	0.700	0.050
0.90011	2020	0.950	1.000	0.650	0.650	0.800	0.050
0.90016	1035	0.950	1.000	0.650	0.650	0.900	0.050

* Power was computed using the normal approximation method.

These scenarios require large sample sizes.

Non-Inferiority Tests for the Ratio of Two Correlated Proportions

Example 3 – Validation using Nam and Blackwelder (2002)

Nam and Blackwelder (2002) give an example in which P_s is 0.80, P_{10} is 0.05, R_1 is 1.00, R_0 is 0.80, the significance level is 0.05, and the power is 80%. From their Table III, the sample size is 34. Note that their calculations use the approximate formula.

Setup

This section presents the values of each of the parameters needed to run this example. First, from the PASS Home window, load the **Non-Inferiority Tests for the Ratio of Two Correlated Proportions** procedure window by expanding **Proportions**, then **Two Correlated Proportions**, then clicking on **Non-Inferiority**, and then clicking on **Non-Inferiority Tests for the Ratio of Two Correlated Proportions**. You may then make the appropriate entries as listed below, or open **Example 3** by going to the **File** menu and choosing **Open Example Template**.

Option	Value
Design Tab	
Find	Sample Size
Power Calculation Method.....	Normal Approximation
Power.....	0.80
Alpha.....	0.05
Rni (Non-Inferiority Ratio).....	0.80
Ra (Actual Ratio)	1.0
P_s (Standard Proportion)	0.80
Nuisance Parameter Type.....	P10 (% +Trt -Std)
Nuisance Parameter Value.....	0.05

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results

Numeric Results							
Hypotheses: $H_0: P_t/P_s \leq R_0$ vs. $H_1: P_t/P_s > R_0$							
	Sample Size	Non-Inf. Ratio	Actual Ratio	Treatment Proportion	Standard Proportion	Nuisance Parameter	
Power	N	R_0	R_1	P_t	P_s	P_{10}	Alpha
0.80050	34	0.800	1.000	0.800	0.800	0.050	0.050
* Power was computed using the normal approximation method.							

The calculated sample size of 34 matches the results of Nam and Blackwelder (2002).