

Chapter 586

Studentized Range Tests

Introduction

This procedure computes power and sample size of tests of whether the means of two or more groups which are analyzed using a studentized range test are different. This is a two-sided test.

Methodology for testing equality among three or more groups has received little attention. An article by Shieh (2018) gives results for two competing test procedures: The F-test and the studentized range test. Results for the F-test are available in **PASS** in another procedure. This procedure provides power and sample size results the studentized range test.

While the F-test is by far the most commonly used method for testing the equality of two or more means, Shieh (2018) showed that neither test is always optimal. In fact, the studentized range test is more powerful when the actual range is close to zero.

Technical Details for the Studentized Range Test

Suppose G groups each have a normal distribution and with means $\mu_1, \mu_2, \dots, \mu_G$ and common variance σ^2 . Let $N_1, N_2, \dots, N_G = N_i$ denote the common sample size of all groups and let N denote the total sample size. In this case of equal group sizes, $N = GN_i$. The multigroup test problem requires one to show that the means are different. Shieh (2018) also considered whether the difference between the minimum and maximum means (the range of the means) is sufficiently small so that the differences among the means can be regarded as of no practical importance. These results are available in **PASS** in a companion procedure.

The One-Way Model

Consider the usual one-way fixed-effects model

$$Y_{gj} = \mu_g + \varepsilon_{gj}$$

where Y_{gj} is response, μ_g are the treatment means, and ε_{gj} are the independent, normally distributed error with zero mean and common variance σ^2 . Here the subscript g indexes the G groups, and the subscript j indexes the N_i subjects in each group.

Cohen (1988) showed that hypotheses about the G means may be obtained using either the variance of the means in terms of the F -test or their range in terms of the studentized range.

Studentized Range Tests

Equality Hypothesis

The hypothesis of mean equality is

$$H_0: \frac{\delta}{\sigma} \leq 0 \text{ versus } H_1: \frac{\delta}{\sigma} > 0$$

where $\delta = \mu_{Max} - \mu_{Min}$ represents the range.

Studentized Range Statistic

The studentized range statistic is defined as follows

$$Q = \frac{\left[\max_{g=1 \text{ to } G} (\bar{Y}_g) - \min_{g=1 \text{ to } G} (\bar{Y}_g) \right] \sqrt{N_i}}{S}$$

where \bar{Y}_g are the sample means and S is the sample variance.

It turns out that the distribution of Q is a function of the pairwise mean differences $\mu_g - \mu_h$, not just the range (the maximum of these differences).

The cumulative distribution function, from which the power can be computed, is given by

$$\Theta(q) = P\{Q \leq q\} = E_K \left\{ \sum_{g=1}^G E_{Z_g} \left[\prod_{\substack{h=1 \\ h \neq g}}^G (\Phi\{Z_g + \delta_{gh}\sqrt{N_i}\} - \Phi\{Z_g + \delta_{gh}\sqrt{N_i} - q\sqrt{K/(N-G)}\}) \right] \right\}$$

where $\delta_{gh} = \mu_g - \mu_h$, K is a chi-square random variable with $N - G$ degrees of freedom, $\Phi\{z\}$ is the CDF of a standard normal distribution, Z_g are independent standard normal random variables, $E_K\{x\}$ is the expectation with respect to K , and $E_{Z_g}\{x\}$ is the expectation with respect to Z_g .

Note that the critical value is based on the set of group means. When testing mean equality, this set of means is just

$$\{\mu_1, \dots, \mu_G\} = \{0, \dots, 0\}$$

If a sample size is desired, it can be determined using a standard binary search algorithm.

Studentized Range Tests

Example 1 – Finding Power

An experiment is being designed to test whether the range of the maximum difference among four group means is greater than zero. The hypothesis test will use studentized range test at a significance level of 0.05. Previous studies have shown a standard deviation of 2. Power calculations assume that the actual range is 2.

To better understand the relationship between power and sample size, the researcher wants to compute the power for several group sample sizes between 10 and 50. The sample sizes will be equal across all groups.

Setup

This section presents the values of each of the parameters needed to run this example. First, from the PASS Home window, load the procedure window. You may then make the appropriate entries as listed below, or open **Example 1** by going to the **File** menu and choosing **Open Example Template**.

<u>Option</u>	<u>Value</u>
Design Tab	
Solve For	Power
Alpha.....	0.05
G (Number of Groups).....	4
Ni (Sample Size Per Group)	10 20 30 40 50
μ 1's Input Type	Enter Range of Means H1
δ 1 (Range of Means H1)	2
σ (Standard Deviation)	2

Annotated Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results

Numeric Results

Number of Groups 4

	Total Sample Size	Sample Size Per Group	H1 Range of Means	Std Dev	Alpha
Power	N	Ni	δ 1	σ	
0.3996	40	10	2	2	0.05
0.7406	80	20	2	2	0.05
0.9105	120	30	2	2	0.05
0.9736	160	40	2	2	0.05
0.9931	200	50	2	2	0.05

References

- Shieh, G. 2018. 'On Detecting a Minimal Important Difference among Standardized Means'. Current Psychology, Vol 37, Pages 640-647. Doi: 10.1007/s12144-016-9549-5
- Cohen, Jacob. 1988. Statistical Power Analysis for the Behavioral Sciences. Lawrence Erlbaum Associates. Hillsdale, New Jersey

Report Definitions


- Power is the probability of rejecting a false null hypothesis in favor of the alternative hypothesis.
- Total Sample Size N is the total number of subjects in the study.
- Sample Size Per Group Ni is the number of subjects sampled per group.
- H1 Range of Means δ 1 is the range of the group means assumed by the alternative hypothesis. It is the value at which the power is computed.
- Std Dev σ is the standard deviation of the responses for all groups.
- Alpha is the significance level of the test: the probability of rejecting the null hypothesis when it is actually true.

Studentized Range Tests

Summary Statements

In a one-way study, a sample of 40 subjects, divided evenly among 4 groups, achieves a power of 40%. This power assumes a studentized range test at a significance level of 0.05. The group subject counts are 10. The group means under the null hypothesis are equal. The range at which the power is computed is 2. The standard deviation of all groups is 2.

This report shows the numeric results of this power study.

Chart Section**Chart Section**

This plot gives a visual presentation of the results in the Numeric Report.

Studentized Range Tests

Example 2 – Finding the Sample Size Necessary to Reject

Continuing with the last example, we will determine how large the sample size would need to have been for $\alpha = 0.05$ and power = 0.80 or 0.9.

Setup

This section presents the values of each of the parameters needed to run this example. First, from the PASS Home window, load the procedure window. You may then make the appropriate entries as listed below, or open

Example 2 by going to the **File** menu and choosing **Open Example Template**.

<u>Option</u>	<u>Value</u>
Design Tab	
Solve For	Sample Size
Power.....	0.8 0.9
Alpha.....	0.05
G (Number of Groups)	4
μ_1 's Input Type	Enter Range of Means H1
δ_1 (Range of Means H1)	2
σ (Standard Deviation)	2

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results

Numeric Results						
Number of Groups 4						
	Total Sample Size N	Sample Size Per Group Ni	H1 Range of Means δ_1	Std Dev σ	Alpha	
Power	92	23	2	2	0.05	
	120	30	2	2	0.05	

This report shows the necessary sample sizes for achieving powers of 0.8 and 0.9.

Studentized Range Tests

Example 3 – Validation using Shieh (2018)

Shieh (2018) page 644 presents an example in which $\alpha = 0.05$, $G = 3$, $\sigma = 3.189$, means under alternative hypothesis are $\{7.77, 9.77, 6.68\}$, and power = 0.8. The sample size was stated as being 22 per group for a total of 66.

Setup

This section presents the values of each of the parameters needed to run this example. First, from the PASS Home window, load the procedure window. You may then make the appropriate entries as listed below, or open **Example 3** by going to the **File** menu and choosing **Open Example Template**.

<u>Option</u>	<u>Value</u>
Design Tab	
Solve For	Sample Size
Power.....	0.8
Alpha.....	0.05
G (Number of Groups).....	3
μ_i 's Input Type	Enter $\mu_{11}, \mu_{12}, \dots, \mu_{1G}$
$\mu_{11}, \mu_{12}, \dots, \mu_{1G}$	7.77 9.77 6.68
σ (Standard Deviation)	3.189

Output

Click the Calculate button to perform the calculations and generate the following output.

Numeric Results

Numeric Results						
Number of Groups 3						
	Total Sample Size	Sample Size Per Group	H1 Group Means Set	H1 Range of Means	Std Dev	Alpha
Power	N	Ni	μ_{1i}	δ_1	σ	
0.8187	66	22	$\mu_{1i(1)}$	3.09	3.189	0.05
Name Values						
$\mu_{1i(1)}$	7.77, 9.77, 6.68					

PASS also found $N_i = 22$ and $N = 66$.