

## Chapter 448

# Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance

---

## Introduction

This procedure computes power and sample size for *superiority by a margin* tests in two-sample designs in which the outcome is a continuous normal random variable and the variances of the two groups (populations) are assumed to be equal. Measurements are made on individuals that have been randomly assigned to one of two groups. This is sometimes referred to as a *parallel-groups* design. This design is used in situations such as the comparison of the income level of two regions, the nitrogen content of two lakes, or the effectiveness of two drugs.

The details of sample size calculation for the two-sample design are presented in the Two-Sample T-Tests Assuming Equal Variance chapter and they will not be duplicated here. This chapter only discusses those changes necessary for superiority tests. Sample size formulas for superiority tests of two means are presented in Chow et al. (2018) pages 50-51.

---

## The Statistical Hypotheses

Remember that in the usual *t*-test setting, the null ( $H_0$ ) and alternative ( $H_1$ ) hypotheses for one-sided tests are defined as

$$H_0: \mu_1 - \mu_2 \leq \delta_0 \quad \text{versus} \quad H_1: \mu_1 - \mu_2 > \delta_0$$

or equivalently

$$H_0: \delta \leq \delta_0 \quad \text{versus} \quad H_1: \delta > \delta_0.$$

Rejecting this test implies that the mean difference is larger than the value  $\delta_0$ . This test is called an *upper-tailed test* because it is rejected in samples in which the difference between the sample means is larger than  $\delta_0$ .

Following is an example of a *lower-tailed test*.

$$H_0: \mu_1 - \mu_2 \geq \delta_0 \quad \text{versus} \quad H_1: \mu_1 - \mu_2 < \delta_0$$

or equivalently

$$H_0: \delta \geq \delta_0 \quad \text{versus} \quad H_1: \delta < \delta_0.$$

## Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance

*Superiority by a Margin* tests are special cases of the above directional tests. It will be convenient to adopt the following specialized notation for the discussion of these tests.

<u>Parameter</u>	<u>PASS Input/Output</u>	<u>Interpretation</u>
$\mu_1$	Not used	<i>Mean</i> of population 1. Population 1 is assumed to consist of those who have received the new treatment.
$\mu_2$	Not used	<i>Mean</i> of population 2. Population 2 is assumed to consist of those who have received the reference treatment.
$M_S$	SM	<i>Margin of superiority</i> . This is a tolerance value that defines the magnitude of difference that is not of practical importance. This may be thought of as the smallest difference from the reference value that is considered to be of practical significance. This value is assumed to be a positive number.
$\delta$	$\delta$	<i>Actual difference</i> . This is the value of $\mu_1 - \mu_2$ , the difference between the means. This is the value at which the power is calculated.

Note that the actual values of  $\mu_1$  and  $\mu_2$  are not needed. Only their difference is needed for power and sample size calculations.

---

### Superiority by a Margin Tests

A *superiority by a margin test* tests that the treatment mean is better than the reference mean by more than the superiority margin. The actual direction of the hypothesis depends on the response variable being studied.

#### Case 1: High Values Good

In this case, higher values are better. The hypotheses are arranged so that rejecting the null hypothesis implies that the treatment mean is greater than the reference mean by at least the margin of superiority. The value of  $\delta$  at which power is calculated must be greater than  $|M_S|$ . The null and alternative hypotheses with  $\delta_0 = |M_S|$  are

$$H_0: \mu_1 \leq \mu_2 + |M_S| \quad \text{versus} \quad H_1: \mu_1 > \mu_2 + |M_S|$$

$$H_0: \mu_1 - \mu_2 \leq |M_S| \quad \text{versus} \quad H_1: \mu_1 - \mu_2 > |M_S|$$

$$H_0: \delta \leq |M_S| \quad \text{versus} \quad H_1: \delta > |M_S|$$

#### Case 2: High Values Bad

In this case, higher values are worse. The hypotheses are arranged so that rejecting the null hypothesis implies that the treatment mean is less than the reference mean by at least the margin of superiority. The value of  $\delta$  at which power is calculated must be less than  $-|M_S|$ . The null and alternative hypotheses with  $\delta_0 = -|M_S|$  are

$$H_0: \mu_1 \geq \mu_2 - |M_S| \quad \text{versus} \quad H_1: \mu_1 < \mu_2 - |M_S|$$

$$H_0: \mu_1 - \mu_2 \geq -|M_S| \quad \text{versus} \quad H_1: \mu_1 - \mu_2 < -|M_S|$$

$$H_0: \delta \geq -|M_S| \quad \text{versus} \quad H_1: \delta < -|M_S|$$

## Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance

**Example**

A superiority test example will set the stage for the discussion of the terminology that follows. Suppose that a test is to be conducted to determine if a new cancer treatment substantially improves mean bone density. The adjusted mean bone density (AMBD) in the population of interest is 0.002300 gm/cm with a standard deviation of 0.000300 gm/cm. Clinicians decide that if the treatment increases AMBD by more than 5% (0.000115 gm/cm), it provides a significant health benefit.

The hypothesis of interest is whether the mean AMBD in the treated group is more than 0.000115 above that of the reference group. The statistical test will be set up so that if the null hypothesis is rejected, the conclusion will be that the new treatment is superior. The value 0.000115 gm/cm is called the *margin of superiority*.

**Two-Sample Equal-Variance T-Test Statistic**

Under the null hypothesis, this test assumes that the two groups of data are simple random samples from a single population of normally-distributed values that all have the same mean and variance. This assumption implies that the data are continuous, and their distribution is symmetric. The calculation of the test statistic for the case when higher response values are better is as follows.

$$t_{df} = \frac{(\bar{X}_1 - \bar{X}_2) - \delta_0}{S_{\bar{X}_1 - \bar{X}_2}}$$

where

$$\bar{X}_k = \frac{\sum_{i=1}^{n_k} X_{ki}}{n_k},$$

$$s_k = \sqrt{\left( \frac{\sum_{i=1}^{n_k} (X_{ki} - \bar{X}_k)^2}{(n_k - 1)} \right)},$$

$$S_{\bar{X}_1 - \bar{X}_2} = \sqrt{\frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2} \left( \frac{1}{n_1} + \frac{1}{n_2} \right)},$$

$$df = n_1 + n_2 - 2,$$

and  $\delta_0$  is the value of the difference hypothesized by the null hypothesis which depends on the magnitude and sign of  $M_S$ .

The null hypothesis is rejected if the computed p-value is less than a specified level (usually 0.05). Otherwise, no conclusion can be reached.

## Computing the Power

When  $\sigma_1 = \sigma_2 = \sigma$ , the power of the equal-variance  $t$ -test is calculated as follows.

1. Find  $t_\alpha$  such that  $1 - T_{df}(t_\alpha) = \alpha$ , where  $T_{df}(x)$  is the area to the left of  $x$  under a central- $t$  distribution with degrees of freedom,  $df = n_1 + n_2 - 2$ .
2. Calculate:  $\sigma_{\bar{x}} = \sigma \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$ .
3. Calculate the noncentrality parameter:  $\lambda = \frac{\delta - \delta_0}{\sigma_{\bar{x}}}$ .
4. Calculate:  $Power = 1 - T'_{df,\lambda}(t_\alpha)$ , where  $T'_{df,\lambda}(x)$  is the area to the left of  $x$  under a noncentral- $t$  distribution with degrees of freedom,  $df = n_1 + n_2 - 2$ , and noncentrality parameter,  $\lambda$ .

When solving for something other than power, PASS uses this same power calculation formulation, but performs a search to determine that parameter.

## Procedure Options

This section describes the options that are specific to this procedure. These are located on the Design tab. For more information about the options of other tabs, go to the Procedure Window chapter.

### Design Tab

The Design tab contains most of the parameters and options that you will be concerned with.

#### Solve For

##### Solve For

This option specifies the parameter to be calculated from the values of the other parameters.

Select *Sample Size* when you want to determine the sample size needed to achieve a given power and alpha.

Select *Power* when you want to calculate the power of an experiment that has already been run.

#### Test

##### Higher Means Are

This option defines whether higher values of the response variable are to be considered better or worse. The choice here determines the direction of the superiority test.

- **Better (H1:  $\delta > SM$ )**

If higher means are Better, the null hypothesis is  $H_0: \delta \leq SM$ , and the alternative hypothesis is  $H_1: \delta > SM$ .

- **Worse (H1:  $\delta < -SM$ )**

If higher means are Worse, the null hypothesis is  $H_0: \delta \geq -SM$ , and the alternative hypothesis is  $H_1: \delta < -SM$ .

## Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance

### Power and Alpha

#### Power

This option specifies one or more values for power. Power is the probability of rejecting a false null hypothesis, and is equal to one minus Beta. Beta is the probability of a type-II error, which occurs when a false null hypothesis is not rejected. In this procedure, a type-II error occurs when you fail to reject the null hypothesis of inferiority when the null hypothesis should be rejected.

Values must be between zero and one. Historically, the value of 0.80 (Beta = 0.20) was used for power. Now, 0.90 (Beta = 0.10) is also commonly used.

A single value may be entered here or a range of values such as *0.8 to 0.95 by 0.05* may be entered.

#### Alpha

This option specifies one or more values for the probability of a type-I error. A type-I error occurs when a true null hypothesis is rejected. In this procedure, a type-I error occurs when you reject the null hypothesis of inferiority when in fact the mean is not non-inferior.

Values must be between zero and one. Historically, the value of 0.05 has been used for alpha. This means that about one test in twenty will falsely reject the null hypothesis. You should pick a value for alpha that represents the risk of a type-I error you are willing to take in your experimental situation.

You may enter a range of values such as *0.01 0.05 0.10* or *0.01 to 0.10 by 0.01*.

### Sample Size (When Solving for Sample Size)

#### Group Allocation

Select the option that describes the constraints on  $N1$  or  $N2$  or both.

The options are

- **Equal ( $N1 = N2$ )**

This selection is used when you wish to have equal sample sizes in each group. Since you are solving for both sample sizes at once, no additional sample size parameters need to be entered.

- **Enter  $N2$ , solve for  $N1$**

Select this option when you wish to fix  $N2$  at some value (or values), and then solve only for  $N1$ . Please note that for some values of  $N2$ , there may not be a value of  $N1$  that is large enough to obtain the desired power.

- **Enter  $R = N2/N1$ , solve for  $N1$  and  $N2$**

For this choice, you set a value for the ratio of  $N2$  to  $N1$ , and then PASS determines the needed  $N1$  and  $N2$ , with this ratio, to obtain the desired power. An equivalent representation of the ratio,  $R$ , is

$$N2 = R * N1.$$

- **Enter percentage in Group 1, solve for  $N1$  and  $N2$**

For this choice, you set a value for the percentage of the total sample size that is in Group 1, and then PASS determines the needed  $N1$  and  $N2$  with this percentage to obtain the desired power.

#### $N2$ (Sample Size, Group 2)

*This option is displayed if Group Allocation = "Enter  $N2$ , solve for  $N1$ "*

$N2$  is the number of items or individuals sampled from the Group 2 population.

$N2$  must be  $\geq 2$ . You can enter a single value or a series of values.

## Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance

### R (Group Sample Size Ratio)

*This option is displayed only if Group Allocation = "Enter R = N2/N1, solve for N1 and N2."*

R is the ratio of N2 to N1. That is,

$$R = N2 / N1.$$

Use this value to fix the ratio of N2 to N1 while solving for N1 and N2. Only sample size combinations with this ratio are considered.

N2 is related to N1 by the formula:

$$N2 = [R \times N1],$$

where the value  $[Y]$  is the next integer  $\geq Y$ .

For example, setting  $R = 2.0$  results in a Group 2 sample size that is double the sample size in Group 1 (e.g.,  $N1 = 10$  and  $N2 = 20$ , or  $N1 = 50$  and  $N2 = 100$ ).

R must be greater than 0. If  $R < 1$ , then N2 will be less than N1; if  $R > 1$ , then N2 will be greater than N1. You can enter a single or a series of values.

### Percent in Group 1

*This option is displayed only if Group Allocation = "Enter percentage in Group 1, solve for N1 and N2."*

Use this value to fix the percentage of the total sample size allocated to Group 1 while solving for N1 and N2. Only sample size combinations with this Group 1 percentage are considered. Small variations from the specified percentage may occur due to the discrete nature of sample sizes.

The Percent in Group 1 must be greater than 0 and less than 100. You can enter a single or a series of values.

---

## Sample Size (When Not Solving for Sample Size)

### Group Allocation

Select the option that describes how individuals in the study will be allocated to Group 1 and to Group 2.

The options are

- **Equal (N1 = N2)**  
This selection is used when you wish to have equal sample sizes in each group. A single per group sample size will be entered.
- **Enter N1 and N2 individually**  
This choice permits you to enter different values for N1 and N2.
- **Enter N1 and R, where N2 = R \* N1**  
Choose this option to specify a value (or values) for N1, and obtain N2 as a ratio (multiple) of N1.
- **Enter total sample size and percentage in Group 1**  
Choose this option to specify a value (or values) for the total sample size (N), obtain N1 as a percentage of N, and then N2 as  $N - N1$ .

## Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance

### Sample Size Per Group

*This option is displayed only if Group Allocation = "Equal ( $N1 = N2$ )."*

The Sample Size Per Group is the number of items or individuals sampled from each of the Group 1 and Group 2 populations. Since the sample sizes are the same in each group, this value is the value for  $N1$ , and also the value for  $N2$ .

The Sample Size Per Group must be  $\geq 2$ . You can enter a single value or a series of values.

### N1 (Sample Size, Group 1)

*This option is displayed if Group Allocation = "Enter  $N1$  and  $N2$  individually" or "Enter  $N1$  and  $R$ , where  $N2 = R * N1$ ."*

$N1$  is the number of items or individuals sampled from the Group 1 population.

$N1$  must be  $\geq 2$ . You can enter a single value or a series of values.

### N2 (Sample Size, Group 2)

*This option is displayed only if Group Allocation = "Enter  $N1$  and  $N2$  individually."*

$N2$  is the number of items or individuals sampled from the Group 2 population.

$N2$  must be  $\geq 2$ . You can enter a single value or a series of values.

### R (Group Sample Size Ratio)

*This option is displayed only if Group Allocation = "Enter  $N1$  and  $R$ , where  $N2 = R * N1$ ."*

$R$  is the ratio of  $N2$  to  $N1$ . That is,

$$R = N2/N1$$

Use this value to obtain  $N2$  as a multiple (or proportion) of  $N1$ .

$N2$  is calculated from  $N1$  using the formula:

$$N2 = [R \times N1],$$

where the value  $[Y]$  is the next integer  $\geq Y$ .

For example, setting  $R = 2.0$  results in a Group 2 sample size that is double the sample size in Group 1.

$R$  must be greater than 0. If  $R < 1$ , then  $N2$  will be less than  $N1$ ; if  $R > 1$ , then  $N2$  will be greater than  $N1$ . You can enter a single value or a series of values.

### Total Sample Size (N)

*This option is displayed only if Group Allocation = "Enter total sample size and percentage in Group 1."*

This is the total sample size, or the sum of the two group sample sizes. This value, along with the percentage of the total sample size in Group 1, implicitly defines  $N1$  and  $N2$ .

The total sample size must be greater than one, but practically, must be greater than 3, since each group sample size needs to be at least 2.

You can enter a single value or a series of values.

### Percent in Group 1

*This option is displayed only if Group Allocation = "Enter total sample size and percentage in Group 1."*

This value fixes the percentage of the total sample size allocated to Group 1. Small variations from the specified percentage may occur due to the discrete nature of sample sizes.

The Percent in Group 1 must be greater than 0 and less than 100. You can enter a single value or a series of values.

## Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance

---

### Effect Size – Mean Difference

#### **SM (Superiority Margin)**

This is the magnitude of the margin of superiority. It must be entered as a positive number. If a negative value is entered, the absolute value is used.

When higher means are better, this value is the distance above the reference mean that is required to be considered superior. When higher means are worse, this value is the distance below the reference mean that is required to be considered superior.

#### **$\delta$ (Actual Difference to Detect)**

This is the actual difference between the treatment mean and the reference mean at which the power is calculated. When higher means are better,  $\delta > SM$ . When higher means are worse,  $\delta < -SM$ .

---

### Effect Size – Standard Deviation

#### **$\sigma$ (Standard Deviation)**

The standard deviation entered here is the assumed standard deviation for both the Group 1 population and the Group 2 population.  $\sigma$  must be a positive number.

When  $\sigma$  is not known, you must supply an estimate. Press the small ' $\sigma$ ' button to the right to obtain calculation options for estimating the standard deviation.



## Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance

## Example 1 – Power Analysis

Suppose that a test is to be conducted to determine if a new cancer treatment improves bone density. The adjusted mean bone density (AMBD) in the population of interest is 0.002300 gm/cm with a standard deviation of 0.000300 gm/cm. Clinicians decide that if the treatment increases AMBD by more than 5% (0.000115 gm/cm), it generates a significant health benefit. They also want to consider what would happen if the margin of superiority is set to 2.5% (0.0000575 gm/cm).

The analysis will be a superiority test at the 0.025 significance level. Power to be calculated assuming that the new treatment has 7.5% improvement on AMBD. Several sample sizes between 10 and 800 will be analyzed. The researchers want to achieve a power of at least 90%. All numbers have been multiplied by 10000 to make the reports and plots easier to read.

### Setup

This section presents the values of each of the parameters needed to run this example. First, from the PASS Home window, load the **Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance** procedure window by expanding **Means**, then **Two Independent Means**, then clicking on **Superiority by a Margin**, and then clicking on **Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance**. You may then make the appropriate entries as listed below, or open **Example 1** by going to the **File** menu and choosing **Open Example Template**.

<u>Option</u>	<u>Value</u>
<b>Design Tab</b>	
Solve For .....	<b>Power</b>
Higher Means Are .....	<b>Better (H1: <math>\delta &gt; SM</math>)</b>
Alpha .....	<b>0.025</b>
Group Allocation .....	<b>Equal (N1 = N2)</b>
Sample Size Per Group .....	<b>10 50 100 200 300 500 600 800</b>
SM (Superiority Margin) .....	<b>0.575 1.15</b>
$\delta$ (Actual Difference to Detect) .....	<b>1.725</b>
$\sigma$ (Standard Deviation) .....	<b>3</b>

### Annotated Output

Click the Calculate button to perform the calculations and generate the following output.

### Numeric Results and Plots

#### Numeric Results for an Equal-Variance T-Test

$$\delta = \mu_1 - \mu_2 = \mu_T - \mu_R$$

Higher Means are Better

Hypotheses: H0:  $\delta \leq SM$  vs. H1:  $\delta > SM$

Power	N1	N2	N	SM	$\delta$	$\sigma$	Alpha
0.12553	10	10	20	0.575	1.725	3.0	0.025
0.47524	50	50	100	0.575	1.725	3.0	0.025
0.76957	100	100	200	0.575	1.725	3.0	0.025
0.96885	200	200	400	0.575	1.725	3.0	0.025
0.99681	300	300	600	0.575	1.725	3.0	0.025
0.99998	500	500	1000	0.575	1.725	3.0	0.025
1.00000	600	600	1200	0.575	1.725	3.0	0.025

(report continues)

## Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance

### References

Chow, S.C., Shao, J., Wang, H., and Lohknygina, Y. 2018. Sample Size Calculations in Clinical Research, Third Edition. Taylor & Francis/CRC. Boca Raton, Florida.  
 Julious, Steven A. 2004. 'Tutorial in Biostatistics. Sample sizes for clinical trials with Normal data.' Statistics in Medicine, 23:1921-1986.

### Report Definitions

Power is the probability of rejecting a false null hypothesis.

N1 and N2 are the number of items sampled from each population.

$N = N1 + N2$  is the total sample size.

SM is the magnitude of the margin of superiority. Since higher means are better, this value is positive and is the distance above the reference mean that is required to be considered superior.

$\delta = \mu1 - \mu2 = \muT - \muR$  is the difference between the treatment and reference means at which power and sample size calculations are made.


$\sigma$  is the assumed population standard deviation for each of the two groups.

Alpha is the probability of rejecting a true null hypothesis.

### Summary Statements

Group sample sizes of 10 and 10 achieve 13% power to detect superiority using a one-sided, two-sample equal-variance t-test. The margin of superiority is 0.575. The actual difference between the means is assumed to be 1.725. The significance level (alpha) of the test is 0.025. The data are drawn from populations with a standard deviation of 3.0 in both groups.

### Chart Section



The above report shows that for SM = 1.15, the sample size necessary to obtain 90% power is about 600 per group. However, if SM = 0.575, the required sample size is only about 150 per group.

## Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance

## Example 2 – Finding the Sample Size

Continuing with Example 1, the researchers want to know the exact sample size for each value of SM to achieve 90% power.

### Setup

This section presents the values of each of the parameters needed to run this example. First, from the PASS Home window, load the **Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance** procedure window by expanding **Means**, then **Two Independent Means**, then clicking on **Superiority by a Margin**, and then clicking on **Two-Sample T-Tests for Superiority by a Margin Assuming Equal Variance**. You may then make the appropriate entries as listed below, or open **Example 2** by going to the **File** menu and choosing **Open Example Template**.

<u>Option</u>	<u>Value</u>
<b>Design Tab</b>	
Solve For .....	<b>Sample Size</b>
Higher Means Are.....	<b>Better (H1: <math>\delta &gt; SM</math>)</b>
Power.....	<b>0.90</b>
Alpha.....	<b>0.025</b>
Group Allocation .....	<b>Equal (N1 = N2)</b>
SM (Superiority Margin).....	<b>0.575 1.15</b>
$\delta$ (Actual Difference to Detect) .....	<b>1.725</b>
$\sigma$ (Standard Deviation) .....	<b>3</b>

### Output

Click the Calculate button to perform the calculations and generate the following output.

#### Numeric Results

##### Numeric Results for an Equal-Variance T-Test

$$\delta = \mu_1 - \mu_2 = \mu_T - \mu_R$$

Higher Means are Better

Hypotheses:  $H_0: \delta \leq SM$  vs.  $H_1: \delta > SM$

Target Power	Actual Power	N1	N2	N	SM	$\delta$	$\sigma$	Alpha
0.90	0.90004	144	144	288	0.575	1.725	3.0	0.025
0.90	0.90036	573	573	1146	1.150	1.725	3.0	0.025

This report shows the exact sample size requirement for each value of SM.

## Example 3 – Validation

This procedure uses the same mechanics as the Two-Sample T-Tests for Non-Inferiority Assuming Equal Variance procedure. We refer the user to Examples 3 and 4 of Chapter 450 for the validation.